



12-28-01

JULI NOV 2001 21 JUL 2001



| | | |
|--|---|---|
| FORM PTO-1390 U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK SERVICE (REV. 10-95) | | ATTORNEY'S DOCKET NUMBER CM2176 U.S. APPLICATION NO. (if known, see 37 CFR 1.5) 10/019178 |
| TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371 | | |
| INTERNATIONAL APPLICATION NO. PCT/US00/18468 | INTERNATIONAL FILING DATE 06 July 2000 | PRIORITY DATE CLAIMED 08 July 1999 |
| TITLE OF INVENTION Process For Producing Particles of Amine Reaction Products | | |
| APPLICANT(S) FOR DO/EO/US BUSCH, Alfred et al. | | |
| Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information. | | |
| 1. <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371. 2. <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371. 3. <input type="checkbox"/> This express request to begin national examination procedures (35 U.S.C. 371(f) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(l). 4. <input checked="" type="checkbox"/> A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date. 5. <input checked="" type="checkbox"/> A copy of the International Application was filed (35 U.S.C. 371(c)(2)) a. <input type="checkbox"/> is transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> has been transmitted by the International Bureau. c. <input checked="" type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US). 6. <input type="checkbox"/> A translation of the International Application into English (35 U.S.C. 371(c)(2)). 7. <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)) a. <input type="checkbox"/> are transmitted herewith (required only if not transmitted by the International Bureau). b. <input type="checkbox"/> have been transmitted by the International Bureau. c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired. d. <input checked="" type="checkbox"/> have not been made and will not be made. 8. <input type="checkbox"/> A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)). 9. <input checked="" type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)). 10. <input type="checkbox"/> A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)). Items 11. to 16. below concern document(s) or information included: 11. <input type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98. 12. <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included. 13. <input checked="" type="checkbox"/> A FIRST preliminary amendment. <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment. 14. <input type="checkbox"/> A substitute specification. 15. <input checked="" type="checkbox"/> A change of power of attorney and/or address letter. 16. <input type="checkbox"/> Other items or information: | | |
| "Express Mail" mailing label number EL483621789715 Date of Deposit 21 Dec 2001 | | |
| I hereby certify that this paper/fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to The Assistant Commissioner of Patents, Washington, D.C. 20231 | | |
| Administrator Mailing Application Signature <i>Virginia O. Best</i> | | |

1001478-1-2001

| | | | | | |
|--|--------------|---|-----------|---|----|
| U.S. APPLICATION NO. 10/019178 | | INTERNATIONAL APPLICATION NO. PCT/US00/18468 | | ATTORNEY'S DOCKET NUMBER CM2176 | |
| | | | | CALCULATIONS PTO USE ONLY | |
| ENTER APPROPRIATE BASIC FEE AMOUNT = | | | | \$740 | |
| Surcharge of \$130.00 for furnishing the oath or declaration later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(e)). | | | | \$0 | |
| CLAIMS | NUMBER FILED | NUMBER EXTRA | RATE | | |
| Total Claims | 22-20 = | 2 | x \$18.00 | \$36 | |
| Independent Claims | 1-3 = | 0 | x \$84.00 | \$0 | |
| MULTIPLE DEPENDENT CLAIM(S) (if applicable) | | | \$280.00 | \$0 | |
| TOTAL OF ABOVE CALCULATIONS = | | | | \$776 | |
| Processing fee of \$130.00 for furnishing the English translation later than [] 20 [] 30 months from the earliest claimed priority date (37 CFR 1.492(f)). | | | | \$0 | |
| TOTAL NATIONAL FEE = | | | | \$776 | |
| Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28,3.31). \$40.00 per property + | | | | \$0 | |
| TOTAL FEES ENCLOSED = | | | | \$776 | |
| | | | | Amount to be refunded | \$ |
| | | | | charged | \$ |
| <p>a. [] A check in the amount of \$ ____ to cover the above fees is enclosed.</p> <p>b. [x] Please charge my Deposit Account No. <u>16-2480</u> in the amount of \$ <u>776</u> cover the above fees. A duplicate copy of this sheet is enclosed.</p> <p>c. [x] The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>16-2480</u>. A duplicate copy of this sheet is enclosed.</p> <p>NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.</p> <p>SEND ALL CORRESPONDENCE TO:</p> | | | | | |
| R. S. Echler, Patent Agent Customer Number 27752 | | | |  Signature T. David Reed Name 32,931 Registration Number | |

TOTAL \$776

10/019170
531 Rec'd PCT 21 DEC 2001

Case CM2176

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the Matter of:

U.S. National Phase Entry

Under 35 USC § 371 from

the International Application of :

Alfred Busch et al. :

Int'l Application No. PCT/US00/18468 :

Filed in the RO/US on 06 July 2000:

Entitled: Process for Producing Particles of Amine Reaction Product

PRELIMINARY AMENDMENT UNDER 37 CFR § 1.112

Assistant Commissioner for Patents

Washington, D.C. 20231

Dear Sir:

Prior to Examination and computation of the fees for entering the captioned International Application into the U.S. National Phase, please preliminarily amend the above-identified application as follows and consider the following Remarks.

AMENDMENTS

IN THE CLAIMS

Please remove the Claim multiple dependencies as follows.

Please amend Claim 3 to depend from Claim 1.

Please amend Claim 4 to depend from Claim 1.

Please amend Claim 5 to depend from Claim 1.

Please amend Claim 6 to depend from Claim 1.

Please amend Claim 7 to depend from Claim 1.

Please amend Claim 8 to depend from Claim 1.

Please amend Claim 9 to depend from Claim 1.

Please amend Claim 10 to depend from Claim 1.

Please amend Claim 11 to depend from Claim 1.

Please amend Claim 13 to depend from Claim 1.

Please amend Claim 15 to depend from Claim 1.

Please amend Claim 16 to depend from Claim 1.

Please amend Claim 20 to depend from Claim 18.

Please cancel Claim 23 without prejudice.

REMARKS


Claim 23 has been canceled without prejudice. The multiple dependency of Claims 3-11, 13, 15-16, and 20 has been changed.

CONCLUSION

Applicants have made an earnest effort to place the present claims in condition for allowance. WHEREFORE, entry of the amendments provided herewith, and allowance of Claims 1-22, as amended, are respectfully requested.

Respectfully submitted,

By



T. David Reed
Agent for Applicants
Registration No. 32,931

20 December 2001
5299 Spring Grove Avenue
Cincinnati, Ohio 45217-1087
Phone: (513) 627-7025
FAX: (513) 627-6333

1001918 1210
T01211 B4T6T001

PROCESS FOR PRODUCING PARTICLES OF AMINE REACTION PRODUCT

5

Field of the invention

The present invention relates to a process for producing particles of amine reaction product.

10

Background of the invention

Perfumed products are well-known in the art. However, consumer acceptance of such perfumed products like laundry and cleaning products is determined not only by the performance achieved with these products but also by the aesthetics associated therewith. The perfume components are therefore an important aspect of the successful formulation of such commercial products.

20

It is also desired by consumers for treated fabrics to maintain the pleasing fragrance over time. Indeed, perfume additives make such compositions more aesthetically pleasing to the consumer, and in some cases the perfume imparts a pleasant fragrance to fabrics treated therewith. However, the amount of perfume carried-over from an aqueous laundry bath onto fabrics is often marginal and does not last long on the fabric. Furthermore, fragrance materials are often very costly and their inefficient use in laundry and cleaning compositions and ineffective delivery to fabrics results in a very high cost to both consumers and laundry and cleaning manufacturers. Industry, therefore, continues to seek with urgency for more efficient and effective fragrance delivery in laundry and cleaning products, especially for improvement in the provision of long-lasting fragrance to the fabrics.

30

It has recently been found that an amine reaction product of a compound containing a primary amine functional group and an active ketone or aldehyde containing component fulfills such a need. Disclosure of such compounds can be

35

found in recently filed applications EP 98870227.0, EP 98870226.2, EP 99870026.4, and EP 99870025.6, all incorporated herein by reference.

However, a problem now encountered with the use of these compound is their ease of incorporation into fully formulated compositions. Indeed, such amine reaction products are often viscous which renders their incorporation into these fully formulated compositions more cumbersome.

It is therefore an object of the invention to provide amine reaction product in a form suitable for easy incorporation into fully formulated composition.

It has now been found that the mixing of the amine reaction product with a carrier having specific melting point to form a particle fulfills such a need. Depending on the property of the carrier, the obtained particle is thereafter treated to form a coated particle.

Further, it has also been advantageously found that both viscous and non-viscous amine reaction product when processed by the present invention exhibit better deposition and long lasting release than amine reaction product which have not been processed according to this process. Not to be bound by theory, it is believed that the mixing with such carrier acts as a shell around the amine reaction product, thus protecting it from the aggressive wash liquor as well as enhancing its fabric deposition properties.

By "viscous", it is meant a product which has a viscosity higher than 1000 cps. The viscosity is measured on a rheometer, TA Instrument CSL²₁₀₀ at a temperature of 25C with a gap setting of 500 microns.

The carrier for use herein has a melting point, measured at atmospheric pressure, of less than 30°C.

Summary of the invention

The present invention is a process for making particles of amine reaction product of a compound containing a primary and/or secondary amine functional group

and an active ketone or aldehyde containing component, and which comprises the steps of :

- a)-providing the amine reaction product, and
- b)-mixing therewith a carrier having a melting point of less than 30°C.

5 In a preferred embodiment of the invention, the processed amine reaction product is further processed to form a coated particle.

In another aspect of the invention, the processed amine reaction product or coated particle is incorporated in a finished composition.

Detailed description of the invention

Starting materials

1) -Amine reaction product

The amine reaction product for use herein is a product of reaction between a compound containing a primary and/or secondary amine functional group and an active ketone or aldehyde containing component, so called hereinafter "amine reaction product".

A typical disclosure of amine reaction product suitable for use herein can be found in recently filed applications EP 98870227.0, EP 98870226.2, EP 99870026.4, and EP 99870025.6, all incorporated herein by reference.

A-Primary and/or secondary amine

By "primary and/or secondary amine", it is meant a component which carries at least one primary and/or secondary amine and/or amide function.

Preferably, the primary and/or secondary amine compound is also characterized by an Odour Intensity Index of less than that of a 1% solution of methyl anthranilate in dipropylene glycol.

Odour Intensity Index method

By Odour Intensity Index, it meant that the pure chemicals were diluted at 1% in Dipropylene Glycol, odor-free solvent used in perfumery. This percentage is more representative of usage levels. Smelling strips, or so called "blotters", were dipped and presented to the expert panellist for evaluation. Expert panellists are assessors trained for at least six months in odor grading and whose gradings are checked for accuracy and reproducibility versus a reference on an on-going basis. For each amine compound, the panellist was presented two blotters: one reference (Me Anthranilate, unknown from the panellist) and the sample. The panellist was asked to rank both smelling strips on the 0-5 odor intensity scale, 0 being no odor detected, 5 being very strong odor present.

Results:

The following represents Odour Intensity Index of an amine compound suitable for use in the present invention and according to the above procedure. In each case, numbers are arithmetic averages among 5 expert panellists and the results are statistically significantly different at 95% confidence level:

| | |
|----------------------------------|-----|
| Methylantranilate 1% (reference) | 3.4 |
| Ethyl-4-aminobenzoate (EAB) 1% | 0.9 |

A general structure for the primary amine compound of the invention is as follows:

$B-(NH_2)_n$;

wherein B is a carrier material, and n is an index of value of at least 1.

Compounds containing a secondary amine group have a structure similar to the above excepted that the compound comprises one or more -NH- groups instead of -NH₂. Further, the compound structure may also have one or more of both -NH₂ and -NH- groups.

Preferred B carriers are inorganic or organic carriers.

By "inorganic carrier", it is meant carrier which are non-or substantially non carbon based backbones.

Preferred primary and/or secondary amines, among the inorganic carriers, are those selected from mono or polymers or organic-organosilicon copolymers of amino derivatised organo silane, siloxane, silazane, alumane, aluminum siloxane, or aluminum silicate compounds. Typical examples of such carriers are: organosiloxanes with at least one primary amine moiety like the diaminoalkylsiloxane $[H_2NCH_2(CH_3)_2Si]_nO$, or the organoaminosilane $(C_6H_5)_3SiNH_2$ described in: Chemistry and Technology of Silicone, W. Noll, Academic Press Inc. 1998, London, pp 209, 106).

Preferred primary and/or secondary amines, among the organic carriers, are those selected from aminoaryl derivatives, polyamines, amino acids and derivatives thereof, substituted amines and amides, glucamines, dendrimers, polyvinylamines and derivatives thereof, and/or copolymer thereof, alkylene polyamine, polyaminoacid and copolymer thereof, cross-linked polyaminoacids, amino substituted polyvinylalcohol, polyoxyethylene bis amine or bis aminoalkyl, aminoalkyl piperazine and derivatives thereof, bis (amino alkyl) alkyl diamine linear or branched, and mixtures thereof.

Preferred aminoaryl derivatives are the amino-benzene derivatives including the alkyl esters of 4-amino benzoate compounds, and more preferably selected from ethyl-4-amino benzoate, phenylethyl-4-aminobenzoate, phenyl-4-aminobenzoate, 4-amino-N'-(3-aminopropyl)-benzamide, and mixtures thereof.

Polyamines suitable for use in the present invention are polyethyleneimines polymers, poly[oxy(methyl-1,2-ethanediyl)], α -(2-aminomethylethyl)- ω -(2-aminomethyl-ethoxy)- (= C.A.S No. 9046-10-0); poly[oxy(methyl-1,2-ethanediyl)], α -hydro-)- ω -(2-aminomethylethoxy)-, ether with 2-ethyl-2-(hydroxymethyl)-1,3-propanediol (= C.A.S. No. 39423-51-3); commercially available under the tradename Jeffamines T-403, D-230, D-400, D-2000; 2,2',2"-triaminotriethylamine; 2,2'-diamino-diethylamine; 3,3'-diamino-dipropylamine, 1,3 bis aminoethyl-cyclohexane commercially available from Mitsibushi and the C12 Sternamines commercially available from Clariant like the C12

Sternamin(propylenamine)_n with $n=3/4$, and mixtures thereof. Preferred polyamines are polyethyleneimines commercially available under the tradename Lupasol like Lupasol FG (MW 800), G20wfv (MW 1300), PR8515 (MW 2000), WF (MW 25000), FC (MW 800), G20 (MW 1300), G35 (MW 1200), G100 (MW 2000), HF (MW 25000), P (MW 750000), PS (MW 750000), SK (MW 2000000), SNA (MW 1000000).

Preferred amino acids for use herein are selected tyrosine, tryptophane, lysine, glutamic acid, glutamine, aspartic acid, arginine, asparagine, phenylalanine, proline, glycine, serine, histidine, threonine, methionine, and mixture thereof, most preferably selected from tyrosine, tryptophane, and mixture thereof. Preferred amino acid derivatives selected from tyrosine ethylate, glycine methylate, tryptophane ethylate, and mixture thereof.

Preferred substituted amines and amides for use herein are selected from nipecotamide, N-coco-1,3-propenediamine; N-oleyl-1,3-propenediamine; N-(tallow alkyl)-1,3-propenediamine; 1,4-diamino cyclohexane; 1,2-diamino-cyclohexane; 1,12-diaminododecane, and mixtures thereof.

Other primary amine compounds suitable for use herein are the glucamines, preferably selected from 2,3,4,5,6-pentamethoxy-glucamine; 6-acetylglucamine, glucamine, and mixture thereof.

Also preferred compounds are the polyethylenimine and/or polypropylenimine dendrimers and the commercially available Starburst[®] polyamidoamines (PAMAM) dendrimers, generation G0-G10 from Dendritech and the dendrimers Astromols[®], generation 1-5 from DSM being DiAminoButane PolyAmine DAB (PA)_x dendrimers with $x = 2^n \times 4$ and n being generally comprised between 0 and 4.

Still other preferred primary and/or secondary amine containing compounds are amino-functional polymers. Preferred amino-functional polymers for use in the present invention are selected from the polyvinylamines, derivatives thereof, copolymer thereof, alkylene polyamine, polyaminoacids and copolymer thereof,

cross-linked polyaminoacids, amino substituted polyvinylalcohol, polyoxyethylene bis amine or bis aminoalkyl, aminoalkyl piperazine and derivatives, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched (TPTA), and mixtures thereof.

5

Polyamino acid is one suitable and preferred class of amino-functional polymer. Polyaminoacids are compounds which are made up of amino acids or chemically modified amino acids. They can contain alanine, serine, aspartic acid, arginine, valine, threonine, glutamic acid, leucine, cysteine, histidine, lysine, isoleucine, tyrosine, asparagine, methionine, proline, tryptophan, phenylalanine, glutamine, glycine or mixtures thereof. In chemically modified amino acids, the amine or acidic function of the amino acid has reacted with a chemical reagent. This is often done to protect these chemical amine and acid functions of the amino acid in a subsequent reaction or to give special properties to the amino acids, like improved solubility. Examples of such chemical modifications are benzyloxycarbonyl, aminobutyric acid, butyl ester, pyroglutamic acid. More examples of common modifications of amino acids and small amino acid fragments can be found in the Bachem, 1996, Peptides and Biochemicals Catalog.

15

Preferred polyamino acids are polylysines, polyarginine, polyglutamine, polyasparagine, polyhistidine, polytryptophane or mixtures thereof. Most preferred are polylysines or polyamino acids where more than 50% of the amino acids are lysine, since the primary amine function in the side chain of the lysine is the most reactive amine of all amino acids.

20

The preferred polyamino acid has a molecular weight of 500 to 10.000.000, more preferably between 5.000 and 750.000.

The polyamino acid can be cross linked. The cross linking can be obtained for example by condensation of the amine group in the side chain of the amino acid like lysine with the carboxyl function on the amino acid or with protein cross linkers like PEG derivatives. The cross linked polyamino acids still need to have free primary and/or secondary amino groups left for reaction with the active ingredient.

30

35

The preferred cross linked polyamino acid has a molecular weight of 20.000 to 10.000.000, more preferably between 200.000 and 2.000.000.

The polyamino acid or the amino acid can be co-polymerized with other reagents like for instance with acids, amides, acyl chlorides. More specifically with aminocaproic acid, adipic acid, ethylhexanoic acid, caprolactam or mixture thereof. The molar ratio used in these copolymers ranges from 1:1 (reagent/ amino acid (lysine)) to 1:20, more preferably from 1:1 to 1:10.

10 The polyamino acid like polylysine can be partially ethoxylated.

Examples and supply of polyaminoacids containing lysine, arginine, glutamine, asparagine are given in the Bachem 1996, Peptides and Biochemicals catalog.

15 The polyaminoacid can be obtained before reaction with the active ingredient, under a salt form. For example polylysine can be supplied as polylysine hydrobromide. Polylysine hydrobromide is commercially available from Sigma, Applichem, Bachem and Fluka.

20 Examples of suitable amino functional polymers containing at least one primary and/or secondary amine group for the purpose of the present invention are :

- Polyvinylamine with a MW of about 300-2.10E6;
- Polyvinylamine alkoxyated with a MW of about 600, 1200 or 3000 and an ethoxylation degree of 0.5;
- 25 - Polyvinylamine vinylalcohol - molar ratio 2:1, polyvinylaminevinylformamide - molar ratio 1:2 and polyvinylamine vinylformamide-molar ratio 2:1;
- Triethylenetetramine, diethylenetriamine, tetraethylenepentamine;
- Bis-aminopropylpiperazine;
- Polyamino acid (L-lysine / lauric acid in a molar ratio of 10/1), Polyamino acid (L-lysine / aminocaproic acid / adipic acid in a molar ratio of 5/5/1),), Polyamino acid (L-lysine / aminocaproic acid /ethylhexanoic acid in a molar ratio of 5/3/1) Polyamino acid (polylysine-cocaprolactam); Polylysine, Polylysine hydrobromide; cross-linked polylysine,
- 30 - amino substituted polyvinylalcohol with a MW ranging from 400-300,000;
- 35 - polyoxyethylene bis [amine] available from e.g. Sigma;
- polyoxyethylene bis [6-aminoheptyl] available from e.g. Sigma;

- N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched (TPTA); and
- 1,4-bis-(3-aminopropyl) piperazine (BNPP).

The more preferred compounds are selected from ethyl-4-amino benzoate, polyethyleneimine polymers commercially available under the tradename Lupasol like Lupasol FG, G20,wfv, PR8515, WF, FC, G20, G35, G100, HF, P, PS, SK, SNA; the diaminobutane dendrimers Astramol[®], polylysine, cross-linked polylysine, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched; 1,4-bis-(3-aminopropyl) piperazine, and mixtures thereof. Even most preferred compounds are those selected from ethyl-4-amino benzoate, polyethyleneimine polymers commercially available under the tradename Lupasol like Lupasol FG, G20,wfv, PR8515, WF, FC, G20, G35, G100, HF, P, PS, SK, SNA; polylysine, cross-linked polylysine, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched, 1,4-bis-(3-aminopropyl) piperazine, and mixtures thereof.

Advantageously, such most preferred primary and/or secondary amine compounds also provide fabric appearance benefit, in particular colour appearance benefit, thus providing a resulting amine reaction product with the dual properties of both fabric appearance benefit and delayed release of the active. Further, when the primary and/or secondary amine compound has more than one free primary and/or secondary amine group, several different active ingredients (aldehyde and/or ketone) can be linked to the amine compound.

B-Active ketone and/or aldehyde

Preferably, for the above mentioned compounds, by active ketone or active aldehyde, it is meant any chain containing at least 1 carbon atom, preferably at least 5 carbon atoms.

Preferably, the active ketone or active aldehyde is respectively selected from a flavour ketone or aldehyde ingredient, a pharmaceutical ketone or aldehyde active, a biocontrol ketone or aldehyde agent, a perfume ketone or aldehyde component and mixtures thereof; most preferably a perfume ketone and/or aldehyde.

Flavour ingredients include spices, flavor enhancers that contribute to the overall flavour perception.

Pharmaceutical actives include drugs.

Biocontrol agents include biocides, antimicrobials, bactericides, fungicides, algaecides, mildewcides, disinfectants, sanitiser like bleach, antiseptics, insecticides, insect and/or moth repellent, vermicides, plant growth hormones.

Typical antimicrobials include Glutaraldehyde, Cinnamaldehyde, and mixtures thereof. Typical insect and/or moth repellants are perfume ingredients, such as citronellal, citral, N, N diethyl meta toluamide, Rotundial, 8-acetoxycarvotanacenone, and mixtures thereof. Other examples of insect and/or moth repellent for use herein are disclosed in US 4,449,987, 4,693,890, 4,696,676, 4,933,371, 5,030,660, 5,196,200, and "Semio Activity of Flavor and Fragrance molecules on various Insect Species", B.D. Mookherjee et al., published in Bioactive Volatile Compounds from Plants, ASC Symposium Series 525, R. Teranishi, R.G. Buttery, and H. Sugisawa, 1993, pp. 35-48.

A typical disclosure of suitable ketone and/or aldehydes, traditionally used in perfumery, can be found in "perfume and Flavor Chemicals", Vol. I and II, S. Arctander, Allured Publishing, 1994, ISBN 0-931710-35-5.

Perfume ketones components include components having odoriferous properties.

Preferably, for the above mentioned compounds, the perfume ketone is selected from buccoxime; iso jasmone; methyl beta naphthyl ketone; musk indanone; tonalid/musk plus; Alpha-Damascone, Beta-Damascone, Delta-Damascone, Iso-Damascone, Damascenone, Damarose, Methyl-Dihydrojasmonate, Menthone, Carvone, Camphor, Fenchone, Alpha-Ionone, Beta-Ionone, Gamma-Methyl so-called Ionone, Fleuramone, Dihydrojasmone, Cis-Jasmone, Iso-E-Super, Methyl-Cedrenyl-ketone or Methyl- Cedrylone, Acetophenone, Methyl-Acetophenone, Para-Methoxy-Acetophenone, Methyl-Beta-Naphtyl-Ketone, Benzyl-Acetone, Benzophenone, Para-Hydroxy-Phenyl-Butanone, Celery Ketone or Livescone, 6-Isopropyldecahydro-2-naphtone, Dimethyl-Octenone, Freskomenthe, 4-(1-Ethoxyvinyl)-3,3,5,5,-tetramethyl-Cyclohexanone, Methyl-Heptenone, 2-(2-(4-Methyl-3-cyclohexen-1-yl)propyl)-cyclopentanone, 1-(p-Menthen-6(2)-yl)-1-

propanone, 4-(4-Hydroxy-3-methoxyphenyl)-2-butanone, 2-Acetyl-3,3-Dimethyl-Norbornane, 6,7-Dihydro-1,1,2,3,3-Pentamethyl-4(5H)-Indanone, 4-Damascol, Dulcinyll or Cassione, Gelsone, Hexalon, Isocyclemone E, Methyl Cyclocitronone, Methyl-Lavender-Ketone, Orivon, Para-tertiary-Butyl-Cyclohexanone, Verdone, 5 Delphone, Muscone, Neobutenone, Plicatone, Veloutone, 2,4,4,7-Tetramethyl-oct-6-en-3-one, Tetrameran, hedione, and mixtures thereof.

More preferably, for the above mentioned compounds, the preferred ketones are selected from Alpha Damascone, Delta Damascone, Iso Damascone, Carvone, 10 Gamma-Methyl-Ionone, Iso-E-Super, 2,4,4,7-Tetramethyl-oct-6-en-3-one, Benzyl Acetone, Beta Damascone, Damascenone, methyl dihydrojasmonate, methyl cedrylone, hedione, and mixtures thereof.

Perfume aldehyde components include components having odoriferous 15 properties.

Preferably, for the above mentioned compounds, the perfume aldehyde is selected from adoxal; anisic aldehyde; cymal; ethyl vanillin; florhydral; helional; heliotropin; hydroxycitronellal; koavone; lauric aldehyde; lyral; methyl nonyl 20 acetaldehyde; P. T. buccinal; phenyl acetaldehyde; undecylenic aldehyde; vanillin; 2,6,10-trimethyl-9-undecenal, 3-dodecen-1-al, alpha-n-amyl cinnamic aldehyde, 4-methoxybenzaldehyde, benzaldehyde, 3-(4-tert butylphenyl)-propanal, 2-methyl-3-(para-methoxyphenyl) propanal, 2-methyl-4-(2,6,6-trimethyl-2(1)-cyclohexen-1-yl) butanal, 3-phenyl-2-propenal, cis-/trans-3,7-dimethyl-2,6- 25 octadien-1-al, 3,7-dimethyl-6-octen-1-al, [(3,7-dimethyl-6-octenyl)oxy] acetaldehyde, 4-isopropylbenzaldehyde, 1,2,3,4,5,6,7,8-octahydro-8,8-dimethyl-2-naphthaldehyde, 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde, 2-methyl-3-(isopropylphenyl)propanal, 1-decanal; decyl aldehyde, 2,6-dimethyl-5-heptenal, 4-(tricyclo[5.2.1.0(2,6)]-decylidene-8)-butanal, octahydro-4,7-methano-1H- 30 indenecarboxaldehyde, 3-ethoxy-4-hydroxy benzaldehyde, para-ethyl-alpha, alpha-dimethyl hydrocinnamaldehyde, alpha-methyl-3,4-(methylenedioxy)-hydrocinnamaldehyde, 3,4-methylenedioxybenzaldehyde, alpha-n-hexyl cinnamic aldehyde, m-cymene-7-carboxaldehyde, alpha-methyl phenyl acetaldehyde, 7-hydroxy-3,7-dimethyl octanal, Undecenal, 2,4,6-trimethyl-3-cyclohexene-1- 35 carboxaldehyde, 4-(3)(4-methyl-3-pentenyl)-3-cyclohexen-carboxaldehyde, 1-dodecanal, 2,4-dimethyl cyclohexene-3-carboxaldehyde, 4-(4-hydroxy-4-methyl

pentyl)-3-cyclohexene-1-carboxaldehyde, 7-methoxy-3,7-dimethyloctan-1-al, 2-methyl undecanal, 2-methyl decanal, 1-nonanal, 1-octanal, 2,6,10-trimethyl-5,9-undecadienal, 2-methyl-3-(4-tertbutyl)propanal, dihydrocinnamic aldehyde, 1-methyl-4-(4-methyl-3-pentenyl)-3-cyclohexene-1-carboxaldehyde, 5 or 6 methoxyhexahydro-4,7-methanoindan-1 or 2- carboxaldehyde, 3,7-dimethyloctan-1-al, 1-undecanal, 10-undecen-1-al, 4-hydroxy-3-methoxy benzaldehyde, 1-methyl-3-(4-methylpentyl)-3-cyclohexenecarboxaldehyde, 7-hydroxy-3,7-dimethyl-octanal, trans-4-decenal, 2,6-nonadienal, para-tolylacetaldehyde; 4-methylphenylacetaldehyde, 2-methyl-4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-2-butenal, ortho-methoxycinnamic aldehyde, 3,5,6-trimethyl-3-cyclohexene carboxaldehyde, 3,7-dimethyl-2-methylene-6-octenal, phenoxyacetaldehyde, 5,9-dimethyl-4,8-decadienal, peony aldehyde (6,10-dimethyl-3-oxa-5,9-undecadien-1-al), hexahydro-4,7-methanoindan-1-carboxaldehyde, 2-methyl octanal, alpha-methyl-4-(1-methyl ethyl) benzene acetaldehyde, 6,6-dimethyl-2-norpinene-2-propionaldehyde, para methyl phenoxy acetaldehyde, 2-methyl-3-phenyl-2-propen-1-al, 3,5,5-trimethyl hexanal, Hexahydro-8,8-dimethyl-2-naphthaldehyde, 3-propyl-bicyclo[2.2.1]-hept-5-ene-2-carbaldehyde, 9-decenal, 3-methyl-5-phenyl-1-pentanal, methyl nonyl acetaldehyde, 1-p-menthene-q-carboxaldehyde, citral, lilial and mixtures thereof.

Most preferred aldehydes are selected from citral, 1-decanal, benzaldehyde, florhydral, 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde; cis/trans-3,7-dimethyl-2,6-octadien-1-al; heliotropin; 2,4,6-trimethyl-3-cyclohexene-1-carboxaldehyde; 2,6-nonadienal; alpha-n-amyl cinnamic aldehyde, alpha-n-hexyl cinnamic aldehyde, P.T. Bucinal, lylal, cymal, methyl nonyl acetaldehyde, trans-2-nonenal, lilial, trans-2-nonenal, and mixture thereof.

In the above list of perfume ingredients, some are commercial names conventionally known to one skilled in the art, and also includes isomers. Such isomers are also suitable for use in the present invention.

In another embodiment, especially suitable for the purpose of the present invention are the perfume compounds, preferably the perfume ketones or aldehydes, characterised by having a low Odor Detection Threshold. Such Odor Detection Threshold (ODT) should be lower than 1ppm, preferably lower than 10ppb - measured at controlled Gas Chromatography (GC) conditions such as

described here below. This parameter refers to the value commonly used in the perfumery arts and which is the lowest concentration at which significant detection takes place that some odorous material is present, as referred to for example in "Compilation of Odor and Taste Threshold Value Data (ASTM DS 48 A)", edited by F. A. Fazzalari, International Business Machines, Hopwell Junction, NY and in Calkin et al., Perfumery, Practice and Principles, John Willey & Sons, Inc., page 243 et seq (1994). For the purpose of the present invention, the Odor Detection Threshold is measured according to the following method :

The gas chromatograph is characterized to determine the exact volume of material injected by the syringe, the precise split ratio, and the hydrocarbon response using a hydrocarbon standard of known concentration and chain-length distribution. The air flow rate is accurately measured and, assuming the duration of a human inhalation to last 0.02 minutes, the sampled volume is calculated. Since the precise concentration at the detector at any point in time is known, the mass per volume inhaled is known and hence the concentration of material. To determine the ODT of a perfume material, solutions are delivered to the sniff port at the back-calculated concentration. A panelist sniffs the GC effluent and identifies the retention time when odor is noticed. The average over all panelists determines the threshold of noticeability. The necessary amount of analyte is injected onto the column to achieve a certain concentration, such as 10 ppb, at the detector. Typical gas chromatograph parameters for determining odor detection thresholds are listed below.

GC: 5890 Series II with FID detector

7673 Autosampler

Column: J&W Scientific DB-1

Length 30 meters ID 0.25 mm film thickness 1 micron

Method:

Split Injection: 17/1 split ratio

Autosampler: 1.13 microliters per injection

Column Flow: 1.10 mL/minute

Air Flow: 345 mL/minute

Inlet Temp. 245°C

Detector Temp. 285°C

Temperature Information

Initial Temperature: 50°C

Rate: 5C/minute

Final Temperature: 280°C

Final Time: 6 minutes

Leading assumptions: 0.02 minutes per sniff

GC air adds to sample dilution

5

Examples of such preferred perfume components are those selected from : 2-methyl-2-(para-iso-propylphenyl)-propionaldehyde, 1-(2,6,6-trimethyl-2-cyclohexan-1-yl)-2-buten-1-one and/or para-methoxy-acetophenone. Even more preferred are the following compounds having an ODT \leq 10ppb measured with the method described above : undecylenic aldehyde, undecalactone gamma, heliotropin, dodecalactone gamma, p-anisic aldehyde, para hydroxy-phenyl-butanone, cymal, benzyl acetone, ionone alpha, p.t.bucinal, damascenone, ionone beta and methyl-nonyl ketone.

10

15

Typically the level of active is of from 10 to 90%, preferably from 30 to 85%, more preferably from 45 to 80% by weight of the amine reaction product.

20

Preferred amine reaction products are those which have a Dry Surface Odour Index as per given in co-pending application EP 98870155.3 given at page 29, line 26 to page 32 line 29, in which the specified unperfumed base for fabric surfaces and hard surfaces are respectively as follow:

| Composition for fabric surface test | % by weight |
|-------------------------------------|-------------|
| LAS | 16 |
| NaSKS-6 | 6 |
| PB1 | 8 |
| TAED | 2.4 |
| Carbonate | 1 |
| Sodium Carbonate | 1 |
| HEDP | 0.4 |
| SRP1 | 0.2 |
| Photobleach | 0.013 |
| Citric acid | 1.0 |
| Protease | 0.3 |
| Lipase | 0.1 |
| Cellulase | 0.1 |

| | |
|--------------------------------|-----|
| Amylase | 0.3 |
| Zeolite | 3.0 |
| TFAA | 3.0 |
| QAS1 | 2.5 |
| Silicone antifoam | 1.0 |
| Misc/minors to balance to 100% | |

| Composition for hard surface test | % by weight |
|--|-------------|
| C12-14 EO 21 | 2 |
| C12-14 EO 5 | 2.5 |
| C9-11 EO 5 | 2.5 |
| LAS | 0.8 |
| Na ₂ CO ₃ | 0.2 |
| Citric acid | 0.8 |
| Caustic acid | 0.5 |
| Fatty acid | 0.5 |
| SCS | 1.5 |
| Water & Misc/Minors to balance to 100% | |

- Most preferred amine reaction products are those resulting from the reaction of polyethyleneimine polymer like Lupasol polymers, BNPP, or TPTA with one or more of the following Alpha Damascone, Delta Damascone, Carvone, Gamma-Methyl-Ionone, Hedione, Florhydral, Lilial, Heliotropine, and 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde. Still other preferred amine reaction products are those resulting from the reaction of Astramol Dendrimers with Carvone as well as those resulting from the reaction of ethyl-4-amino benzoate with one or more of the following 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde, and trans-2-nonenal. Still another preferred amine reaction products are those resulting from the reaction of polylysine with one or more of the following Alpha Damascone, Delta Damascone, Carvone, and 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde.
- Even most preferred amine reaction products are those from the reaction of Lupasol HF with Delta Damascone; LupasolG35 with Alpha Damascone; LupasolG100 with 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde, BNPP or TPTA

with Alpha and Delta Damascone; ethyl-4-amino benzoate with 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde.

Most preferred amine reaction products are those fulfilling the Dry Surface Odor Index as per given in co-pending application EP 98870155.3 given at page 29, line 26 to page 32 line 29.

2)-Carrier

Another essential ingredient of the process invention is a carrier having a melting point of less than 30°C, preferably between minus 150°C and less than 30°C. By means of this carrier, particles of amine reaction product, suitable for spray-on application, will be produced.

Typical of such carrier includes liquid as well as solid carrier material. The liquid carrier can be in any suitable physical form like hydrophobic or hydrophilic form.

As used herein in relation to carrier materials, "hydrophobic" means substantially water insoluble; "hydrophilic" means substantially water soluble. In this regard, "substantially water insoluble" shall refer to a material that is not soluble in distilled (or equivalent) water, at 25.degree. C., at a concentration of 0.2% by weight, and preferably not soluble at 0.1% by weight (calculated on a water plus carrier weight basis). "Substantially water soluble" shall refer to a material that is soluble in distilled (or equivalent) water, at 25.degree. C., at a concentration of 0.2% by weight, and are preferably soluble at 1.0% by weight. The terms "soluble", "solubility" and the like, for purposes hereof, corresponds to the maximum concentration of carrier, as applicable, that can dissolve in water or other solvents to form a homogeneous solution, as is well understood to those skilled in the art.

Typical of materials that are in hydrophobic form include ingredients useful in perfumery. Typical of such ingredients are the perfume raw materials, solvents, and mixtures thereof. Typical hydrophobic raw materials include the ketones or aldehydes like hexyl cinnamic aldehyde, hydrocarbons like Limolene, d-Limonene, esters like Hercolyn D, benzyl salicylate, hexyl salicylate, triethyl citrate, iso propyl myristate, or mixtures thereof, or hydrophobic perfume composition preferably incorporating one or more of these raw materials.

Typical hydrophobic solvents include diethyl phthalate, ethers like butoxypropoxypropanol, and mixtures thereof.

Also suitable for use herein are the finished perfume compositions which have hydrophobic properties.

Typical of materials that are in hydrophilic form include alcohols like methanol, ethanol, dipropylene glycol, water, nonionic surfactants as above described but having a melting point below 30C, or mixtures thereof.

Preferred carrier materials are selected from benzyl salicylate, diethyl phthalate, dipropylene glycol, methanol, ethanol, and mixtures thereof.

Typically when the amine reaction product is only mixed with a carrier but not further processed, the amine reaction product will be present in an amount of from 1 to 75%, preferably 10 to 60%, more preferably 15 to 45% by weight of the processed reaction product. In this instance, the amount of carrier will be sufficient to add up to 100%. Of course, the processed amine reaction product may also contain minors but in quantities which will not exceed the amount of carrier material. Typically the carrier will be present in an amount of from 3 to 95%, preferably from 15 to 90% and most preferably from 45 to 85%, by weight of the processed amine reaction product.

By processed amine reaction product, it is meant the resulting product of the amine reaction product with the carrier material.

Processing of the amine reaction product with the carrier is done by thoroughly mixing the amine reaction product with the carrier. Advantageously, there is no need for additional ingredients to provide a resulting substantially homogenous mixture. This mixing is done at the lowest possible temperature, i.e. just above the melting point of the carrier or at room temperature. The mixing step is carried out until a complete homogeneous mixture is obtained. By "homogeneous", it is meant compositions which have similar appearance to the resulting composition of a 20g of amine reaction product mixed with 20g of methanol or 80g of diethylphthalate for 5 minutes by means of an Ultra Turrax, the temperature of mixing being room temperature.

The obtained processed amine reaction product may then be suitable for spray on applications when use of a liquid carrier is made.

5 Coating agents

When use of a carrier with a melting point of less than 30 °C is made for the mixing with the amine reaction product, it may then be preferred to further process the mixture to form a coated particle like e.g. by adsorption of the mixture onto a solid, preferably porous coating. The coating will advantageously protect the processed amine reaction product from the aggressive chemicals present in wash water.

The resulting coated particles can be in any form which is suitable for incorporation into liquid or powders, preferably powders, such as agglomerate, pellets, tablets, or mixtures thereof.

Suitable coating agents for both solid, including paste, and liquid mixtures are substantially water-soluble solid binder or agglomerating agents.

"Substantially water soluble" shall refer to a material that is soluble in distilled (or equivalent) water, at 25°C., at a concentration of 0.2% by weight, and are preferably soluble at 1.0% by weight.

A "solid" is defined as a material that is a solid at ambient temperatures, and so solid substantially water-soluble binder or agglomerating agent must have a melting point of at least 30°C, and preferably of at least 40°C.

Suitable water-soluble binders or agglomerating agents include the water soluble organic polymeric compounds, water soluble monomeric polycarboxylates, or their acid forms, homo or copolymeric polycarboxylic acids or their salts in which the polycarboxylic acid comprises at least two carboxylic radicals separated from each other by not more than two carbon atoms, carbonates, bicarbonates, borates, phosphates, sulfate salts like sodium and magnesium sulfate, inorganic perhydrate salts including perborate like perborate monohydrate, percarbonate, silicates, starch, cyclodextrin, and mixtures of any of the foregoing.

Suitable organic polymeric compounds suitable as coating agents include cellulose derivatives such as methylcellulose, carboxymethylcellulose, hydroxypropylcellulose and hydroxyethylcellulose, as well as carbohydrates like pectins, and gums. Further compounds are carbohydrates and derivatives such as fructose, xylose, galactose, galacturonic acid or glucose based polymers like inuline, dextran, xyloglucan, pectin or gums

Suitable carboxylates containing one carboxy group include the water soluble salts of lactic acid, glycolic acid and ether derivatives thereof. Polycarboxylates containing two carboxy groups include the water-soluble salts of succinic acid, malonic acid, (ethylenedioxy) diacetic acid, maleic acid, diglycolic acid, tartaric acid, tartronic acid and fumaric acid, as well as the ether carboxylates and the sulfinyl carboxylates. Polycarboxylates containing three carboxy groups include, in particular, water-soluble citrates, aconitrates and citraconates as well as succinate derivatives such as the carboxymethyloxysuccinates described in British Patent No. 1,379,241, lactoxysuccinates described in British Patent No. 1,389,732, and aminosuccinates described in Netherlands Application 7205873, and the oxypolycarboxylate materials such as 2-oxa-1,1,3-propane tricarboxylates described in British Patent No. 1,387,447.

Polycarboxylates containing four carboxy groups include oxydisuccinates disclosed in British Patent No. 1,261,829, 1,1,2,2-ethane tetracarboxylates, 1,1,3,3-propane tetracarboxylates and 1,1,2,3-propane tetracarboxylates. Polycarboxylates containing sulfo substituents include the sulfosuccinate derivatives disclosed in British Patent Nos. 1,398,421 and 1,398,422 and in U.S. Patent No. 3,936,448, and the sulfonated pyrolysed citrates described in British Patent No. 1,439,000.

Alicyclic and heterocyclic polycarboxylates include cyclopentane-cis,cis,cis-tetracarboxylates, cyclopentadienide pentacarboxylates, 2,3,4,5-tetrahydrofuran - cis, cis, cis-tetracarboxylates, 2,5-tetrahydrofuran - cis - dicarboxylates, 2,2,5,5-tetrahydrofuran - tetracarboxylates, 1,2,3,4,5,6-hexane - hexacarboxylates and carboxymethyl derivatives of polyhydric alcohols such as sorbitol, mannitol and xylitol. Aromatic polycarboxylates include mellitic acid, pyromellitic acid and the phthalic acid derivatives disclosed in British Patent No. 1,425,343.

Of the above, the preferred polycarboxylates are hydroxycarboxylates containing up to three carboxy groups per molecule, more particularly citrates.

Borate, as well as builders containing borate-forming materials that can produce borate under detergent storage or wash conditions can also be used but are not preferred at wash conditions less than about 50°C, especially less than about 40°C.

Examples of carbonates are the alkaline earth and alkali metal carbonates, including sodium carbonate and sesqui-carbonate and mixtures thereof with ultra-fine calcium carbonate as disclosed in German Patent Application No. 2,321,001 published on November 15, 1973.


Specific examples of water-soluble phosphates are the alkali metal tripolyphosphates, sodium, potassium and ammonium pyrophosphate, sodium and potassium and ammonium pyrophosphate, sodium and potassium orthophosphate, sodium polymeta/phosphate in which the degree of polymerization ranges from about 6 to 21, and salts of phytic acid.

Suitable silicates include the water soluble sodium silicates with an SiO_2 : Na_2O ratio of from 1.0 to 2.8, with ratios of from 1.6 to 2.4 being preferred, and 2.0 ratio being most preferred. The silicates may be in the form of either the anhydrous salt or a hydrated salt. Sodium silicate with an SiO_2 : Na_2O ratio of 2.0 is the most preferred silicate.

Typical disclosure of cyclodextrin derivatives are disclosed in WO96/05358, U.S. Pat. Nos: 3,426,011, Parmerter et al., issued Feb. 4, 1969; 3,453,257; 3,453,258; 3,453,259; and 3,453,260, all in the names of Parmerter et al., and all issued July 1, 1969; 3,459,731, Gramera et al., issued Aug. 5, 1969; 3,553,191, Parmerter et al., issued Jan. 5, 1971; 3,565,887, Parmerter et al., issued Feb. 23, 1971; 4,535,152, Szejtli et al., issued Aug. 13, 1985; 4,616,008, Hirai et al., issued Oct. 7, 1986; 4,678,598, Ogino et al., issued Jul. 7, 1987; 4,638,058, Brandt et al., issued Jan. 20, 1987; and 4,746,734, Tsuchiyama et al., issued May 24, 1988; all of said patents being incorporated herein by reference.

Although less preferred for use herein because of their lower solubility, partially water soluble coating agents may also be used as coating agent. These compounds are indeed less preferred because during the wash cycle the amine reaction product will still be at least partially coated and therefore can not display its full functionality of long lasting freshness on dry fabrics or hard surfaces. Examples of partially water soluble coating agents include the crystalline layered silicates. Examples of largely water insoluble builders include the sodium aluminosilicates.


Crystalline layered sodium silicates have the general formula



wherein M is sodium or hydrogen, x is a number from 1.9 to 4 and y is a number from 0 to 20. Crystalline layered sodium silicates of this type are disclosed in EP-A-0164514 and methods for their preparation are disclosed in DE-A-3417649 and DE-A-3742043. For the purpose of the present invention, x in the general formula above has a value of 2, 3 or 4 and is preferably 2. The most preferred material is $\delta\text{-Na}_2\text{Si}_2\text{O}_5$, available from Hoechst AG as NaSKS-6.

Suitable aluminosilicate zeolites have the unit cell formula $\text{Na}_z[(\text{AlO}_2)_z(\text{SiO}_2)_y] \cdot x\text{H}_2\text{O}$ wherein z and y are at least 6; the molar ratio of z to y is from 1.0 to 0.5 and x is at least 5, preferably from 7.5 to 276, more preferably from 10 to 264. The aluminosilicate material are in hydrated form and are preferably crystalline, containing from 10% to 28%, more preferably from 18% to 22% water in bound form.

The aluminosilicate ion exchange materials can be naturally occurring materials, but are preferably synthetically derived. Synthetic crystalline aluminosilicate ion exchange materials are available under the designations Zeolite A, Zeolite B, Zeolite P, Zeolite X, Zeolite MAP, Zeolite HS and mixtures thereof. Zeolite A has the formula



wherein x is from 20 to 30, especially 27. Zeolite X has the formula $\text{Na}_{86}[(\text{AlO}_2)_{86}(\text{SiO}_2)_{106}]\cdot 276 \text{ H}_2\text{O}$.

Typically when the amine reaction product is mixed with a carrier and further processed to form a coated particle, the amine reaction product will be present in an amount of from 1 to 75%, preferably 5 to 30%, more preferably 6 to 25% by weight of the processed reaction product in the produced particle.

Typically the coating agent will be present in an amount of from 10% to 95%, preferably from 30 to 90%, more preferably, 50 to 75% by weight of the particle of the processed amine reaction product. In this instance, the amount of carrier will be sufficient to add up to 100%. Of course, the coated particle may also contain minors but in quantities which will not exceed either of the amount of carrier material or coating agent.

Preferred coating materials are selected from carbonate, starch, cyclodextrin, sulfate salts, silicates, and mixtures thereof.

The surface treatment of the particle can be carried out in a number of ways using equipment known in the art and the process may take in batch wise or continuous fashion.

One method for applying the coating material involves agglomeration. Any conventional agglomerator/mixer may be used including, but not limited to pan, rotary drum and vertical blender types. Molten coating compositions may also be applied either by being poured onto, or spray atomized onto a moving bed of the mixture of amine reaction product with carrier.

Another method for applying the coating is to pour the obtained mixture (so-called particle), as herein before described, onto the coating material and agglomerate it in a Braun Mixer. Care is also taken that the temperature during the mixing and/or coating step does not substantially exceed the melting point of the carrier material. For example, 50g of a mixture containing a hydrophobic perfume composition and 25g of the amine reaction product is poured at room temperature into a Braun Mixer containing 200g of starch. The mixing of the ingredients is carried out for about 5 minutes.

Once the coated particle is obtained, it may be useful to remove the carrier. Such removal can be made by conventional techniques. One typical of these technique is vacuum drying. By removal of the carrier, in particular with liquid carrier having a boiling point between 5 and 200°C, the original viscosity of the amine reaction product, i.e. higher viscosity, in the coated particle is restored which as a result gives a better stability of the coated particle towards the wash liquor environment. The coated particle can then be used as is for incorporation into the finished composition.

Accordingly, there is provided a processed amine reaction product as obtainable by the process of the invention.

If desired, the coated particle may also contain one or more additional ingredients like a surfactant for improved solubility or dispersability. Typical of such surfactant are the anionic, nonionic, or cationic surfactants. a cationic, anion or non-ionic surfactant like. Preferably, the weight ratio of such additional ingredient(s) to the coating agent is of up to 1:1.

In another preferred marketing execution, an additional coating on the coated particle can be provided, which depending on the nature of this additional coating will give improved storage stability, flowability and/or improved fabric substantivity of the coated particle. One typical example is polyvinyl alcohol.

Incorporation into finished composition

The finished compositions aspect of the invention, including laundry compositions, hard surface cleaning compositions, personal cleaning compositions, comprises the incorporation of the hereinbefore described processed amine reaction product together with one or more laundry or cleaning ingredient in a finished composition. Laundry compositions of the invention, which also encompass compositions providing color care, are suitable for use in any steps of the domestic treatment, that is as a pre-treatment composition, as a wash additive as a composition suitable for use in the rinse-cycle of the laundry cycle or applied on a dryer-sheet. Obviously, multiple applications can be made such as treating the fabric with a pre-treatment composition of the invention and

also thereafter with a composition of the invention suitable for use in the rinse cycle and/or suitable for use as a dryer-sheet.

5 As mentioned hereinbefore, the incorporation of the processed amine reaction product is conveniently made depending on its end form by either by spraying when in sprayable liquid form, or dry-addition when in coated form.

10 Laundry compositions encompass laundry detergent compositions, including liquid, solid form like powdered, tablets as well as softening compositions including rinse added softening composition as well as dryer added softening compositions.

15 A conventional disclosure of softening ingredients to be used in the softening composition of the invention can be found in EP 98870227.0, incorporated herein by reference.

20 Finished compositions incorporating the processed amine reaction product will normally contain from 0.001 to 25%, more preferably from 0.02 to 10%, and most preferably from 0.05 to 5% of the processed product on a composition weight basis.

Preferably, the finished composition is a detergent composition, more preferably in solid form.

25 In particular, it is preferred that the detergent composition comprises a clay.

Clay

30 The compositions of the invention may preferably contain a clay, preferably present at a level of from 0.05% to 40%, more preferably from 0.5% to 30%, most preferably from 2% to 20% by weight of the composition. For clarity, it is noted that the term clay mineral compound, as used herein, excludes sodium aluminosilicate zeolite builder compounds, which however, may be included in the compositions of the invention as optional components.

35 One preferred clay may be a bentonite clay. Highly preferred are smectite clays, as for example disclosed in the US Patents No.s 3,862,058 3,948,790, 3,954,632

and 4,062,647 and European Patents No.s EP-A-299,575 and EP-A-313,146 all in the name of the Procter and Gamble Company.

The term smectite clays herein includes both the clays in which aluminium oxide is present in a silicate lattice and the clays in which magnesium oxide is present in a silicate lattice. Typical smectite clay compounds include the compounds having the general formula $\text{Al}_2(\text{Si}_2\text{O}_5)_2(\text{OH})_2 \cdot n\text{H}_2\text{O}$ and the compounds having the general formula $\text{Mg}_3(\text{Si}_2\text{O}_5)_2(\text{OH})_2 \cdot n\text{H}_2\text{O}$. Smectite clays tend to adopt an expandable three layer structure.

Specific examples of suitable smectite clays include those selected from the classes of the montmorillonites, hectorites, volchonskoites, nontronites, saponites and sauconites, particularly those having an alkali or alkaline earth metal ion within the crystal lattice structure. Sodium or calcium montmorillonite are particularly preferred.

Suitable smectite clays, particularly montmorillonites, are sold by various suppliers including English China Clays, Laviosa, Georgia Kaolin and Colin Stewart Minerals.

Clays for use herein preferably have a particle dimension of from 10nm to 800nm more preferably from 20nm to 500 nm, most preferably from 50nm to 200 nm.

Particles of the clay mineral compound may be included as components of agglomerate particles containing other detergent compounds. Where present as such components, the term "largest particle dimension" of the clay mineral compound refers to the largest dimension of the clay mineral component as such, and not to the agglomerated particle as a whole.

Substitution of small cations, such as protons, sodium ions, potassium ions, magnesium ions and calcium ions, and of certain organic molecules including those having positively charged functional groups can typically take place within the crystal lattice structure of the smectite clays. A clay may be chosen for its ability to preferentially absorb one cation type, such ability being assessed by measurements of relative ion exchange capacity. The smectite clays suitable herein typically have a cation exchange capacity of at least 50 meq/100g. U.S.

Patent No. 3,954,632 describes a method for measurement of cation exchange capacity.

5 The crystal lattice structure of the clay mineral compounds may have, in a preferred execution, a cationic fabric softening agent substituted therein. Such substituted clays have been termed 'hydrophobically activated' clays. The cationic fabric softening agents are typically present at a weight ratio, cationic fabric softening agent to clay, of from 1:200 to 1:10, preferably from 1:100 to 1:20. Suitable cationic fabric softening agents include the water insoluble tertiary amines or dilong chain amide materials as disclosed in GB-A-1 514 276 and EP-
10 B-0 011 340.

15 A preferred commercially available "hydrophobically activated" clay is a bentonite clay containing approximately 40% by weight of a dimethyl ditallow quaternary ammonium salt sold under the tradename Claytone EM by English China Clays International.

20 In a highly preferred embodiment of the invention, the clay is present in an intimate mixture or in a particle with a humectant and a hydrophobic compound, preferably a wax or oil, such as paraffin oil. Preferred humectants are organic compounds, including propylene glycol, ethylene glycol, dimers or trimers of glycol, most preferably glycerol. The particle is preferably an agglomerate. Alternatively, the particle may be such that the wax or oil and optionally the humectant form an encapsulate on the clay or alternatively, the clay be a
25 encapsulate for the wax or oil and the humectant. It may be preferred that the particle comprises an organic salt or silica or silicate.

30 However, in another embodiment of the invention, the clay is preferably mixed with one or more surfactants and optionally builders and optionally water, in which case the mixture is preferably subsequently dried. Preferably, such a mixture is further processed in a spray-drying method to obtain a spray dried particle comprising the clay.

35 It may be preferred that the flocculating agent is also comprised in the particle or granule comprising the clay.

It may also be preferred that the intimate mixture comprises a chelating agent.

Flocculating agent

5 The compositions of the invention may contain a clay flocculating agent, preferably present at a level of from 0.005% to 10%, more preferably from 0.05% to 5%, most preferably from 0.1% to 2% by weight of the composition.

10 The clay flocculating agent functions such as to bring together the particles of clay compound in the wash solution and hence to aid their deposition onto the surface of the fabrics in the wash. This functional requirement is hence different from that of clay dispersant compounds which are commonly added to laundry detergent compositions to aid the removal of clay soils from fabrics and enable their dispersion within the wash solution.

15 Preferred as clay flocculating agents herein are organic polymeric materials having an average weight of from 100,000 to 10,000,000, preferably from 150,000 to 5,000,000, more preferably from 200,000 to 2,000,000.

20 Suitable organic polymeric materials comprise homopolymers or copolymers containing monomeric units selected from alkylene oxide, particularly ethylene oxide, acrylamide, acrylic acid, vinyl alcohol, vinyl pyrrolidone, and ethylene imine. Homopolymers of, on particular, ethylene oxide, but also acrylamide and acrylic acid are preferred.

25 European Patents No.s EP-A-299,575 and EP-A-313,146 in the name of the Procter and Gamble Company describe preferred organic polymeric clay flocculating agents for use herein.

30 The weight ratio of clay to the flocculating polymer is preferably from 1000:1 to 1:1, more preferably from 500:1 to 1:1, most preferably from 300:1 to 1:1, or even more preferably from 80:1 to 10:1, or in certain applications even from 60:1 to 20:1.


35 Inorganic clay flocculating agents are also suitable herein, typical examples of which include lime and alum.

The flocculating agent is preferably present in a detergent base granule such as a detergent agglomerate, extrudate or spray-dried particle, comprising generally one or more surfactants and builders.

5 Effervescent

Effervescent means may also be optionally used in the compositions of the invention.

10 Effervescency as defined herein means the evolution of bubbles of gas from a liquid, as the result of a chemical reaction between a soluble acid source and an alkali metal carbonate, to produce carbon dioxide gas,



15 Further examples of acid and carbonate sources and other effervescent systems may be found in : (Pharmaceutical Dosage Forms : Tablets Volume 1 Page 287 to 291).

Carbonate salts

20 Suitable alkali and/ or earth alkali inorganic carbonate salts herein include carbonate and hydrogen carbonate of potassium, lithium, sodium, and the like amongst which sodium and potassium carbonate are preferred. Suitable bicarbonates to be used herein include any alkali metal salt of bicarbonate like lithium, sodium, potassium and the like, amongst which sodium and potassium bicarbonate are preferred. However, the choice of carbonate or bicarbonate or
25 mixtures thereof may be made depending on the pH desired in the aqueous medium wherein the granules are dissolved. For example where a relative high pH is desired in the aqueous medium (e.g., above pH 9.5) it may be preferred to use carbonate alone or to use a combination of carbonate and bicarbonate wherein the level of carbonate is higher than the level of bicarbonate. The
30 inorganic alkali and/ or earth alkali carbonate salt of the compositions of the invention comprises preferably a potassium or more preferably a sodium salt of carbonate and/ or bicarbonate. Preferably, the carbonate salt comprises sodium carbonate, optionally also a sodium bicarbonate.

35 The inorganic carbonate salts herein are preferably present at a level of at least 20% by weight of the composition. Preferably they are present at a level of at

least 23% or even 25% or even 30% by weight, preferably up to about 60% by weight or more preferably up to 55% or even 50% by weight.

5 They may be added completely or partially as separate powdered or granular component, as co-granules with other detergent ingredients, for example other salts or surfactants. In solid detergent compositions of the invention, they may also completely or partially be present in detergent granules such as agglomerates or spray dried granules.

10 In one embodiment of the invention, an effervescence source is present, preferably comprising an organic acid, such as carboxylic acids or aminoacids, and a carbonate. Then it may be preferred that part or all of the carbonate salt herein is premixed with the organic acid, and thus present in an separate granular component.

Preferred effervescent source are selected from compressed particles of citric acid and carbonate optionally with a binder; and particle of carbonate, bicarbonate and malic or maleic acid in weight ratios of 4:2:4. The dry add form of citric acid and carbonate are preferably used.

The carbonate may have any particle size. In one embodiment, in particular when the carbonate salt is present in a granule and not as separately added compound, the carbonate salt has preferably a volume median particle size from 5 to 375 microns, whereby preferably at least 60%, preferably at least 70% or even at least 80% or even at least 90% by volume, has a particle size of from 1 to 425 microns. More preferably, the carbon dioxide source has a volume median particle size of 10 to 250, whereby preferably at least 60 %, or even at least 70% or even at least 80% or even at least 90% by volume, has a particle size of from 1 to 375 microns; or even preferably a volume median particle size from 10 to 200 microns, whereby preferably at least 60 %, preferably at least 70% or even at least 80% or even at least 90% by volume, has a particle size of from 1 to 250 microns.

15

In particular when the carbonate salt is added as separate component, so to say 'dry-added' or admixed to the other detergent ingredients, the carbonate may have any particle size, including the above specified particle sizes, but preferably

at least an volume average particle size of 200 microns or even 250 microns or even 300 microns.

- 5 It may be preferred that the carbon dioxide source of the required particle size is obtained by grinding a larger particle size material, optionally followed by selecting the material with the required particle size by any suitable method.

Whilst percarbonate salts may be present in the compositions of the invention as a bleaching agent, they are not included in the carbonate salts as defined herein.

10

The preferred detergent composition, embodiment of the invention, will, also preferably contain a bleach precursor, a source of alkaline hydrogen peroxide necessary to form a peroxyacid bleaching species in the wash solution and preferably will also contain other components conventional in detergent compositions. Thus, preferred detergent compositions will incorporate one or more of surfactants, organic and inorganic builders, soil suspending and anti-redeposition agents, suds suppressors, enzymes, fluorescent whitening agents, photoactivated bleaches, perfumes, colours, and mixtures thereof.

15

20

Typical disclosure of such components can be found in EP-A-0,659,876 and European patent application No. 98870226.2 which are both incorporated herein by reference.

Form of the composition

- 25 The composition of the invention may take a variety of physical form including liquid, gel, foam in either aqueous or non-aqueous form, granular and tablet forms.

30 Still in another aspect of the invention, there is provided a packaged composition comprising the processed product of the invention or composition of the invention. Preferably, the packaged composition is a closed packaging system having a moisture vapour transmission rate of less than $20\text{g/m}^2/24$ hours. Typical disclosure of such a package can be found in WO 98/40464.

- 35 Still another preferred package is a spray dispenser.

Spray Dispenser

The present invention also relates to such compositions incorporated into a spray dispenser to create an article of manufacture that can facilitate treatment of fabric articles and/or surfaces with said compositions containing the amine reaction product and other ingredients (examples are cyclodextrins, polysaccharides, polymers, surfactant, perfume, softener) at a level that is effective, yet is not discernible when dried on the surfaces. The spray dispenser comprises manually activated and non-manual powered (operated) spray means and a container containing the treating composition. Typical disclosure of such spray dispenser can be found in WO 96/04940 page 19 line 21 to page 22 line 27. The articles of manufacture preferably are in association with instructions for use to ensure that the consumer applies sufficient ingredient of the composition to provide the desired benefit. Typical compositions to be dispensed from a sprayer contain a level of amine reaction product of from about 0.01% to about 5%, preferably from about 0.05% to about 2%, more preferably from about 0.1% to about 1%, by weight of the usage composition.

Method of use

Also provided herein is a method for providing a delayed release of an active ketone or aldehyde which comprises the step of contacting the surface to be treated with a compound or composition of the invention, and thereafter contacting the treated surface with a material, preferably an aqueous medium like moisture or any other means susceptible of releasing the perfume from the amine reaction product.

By "surface", it is meant any surface onto which the compound can deposit. Typical examples of such material are fabrics, hard surfaces such as dishware, floors, bathrooms, toilet, kitchen and other surfaces in need of a delayed release of a perfume ketone and/or aldehyde such as that with litter like animal litter. Preferably, the surface is selected from a fabric, a tile, a ceramic; more preferably is a fabric.

By "delayed release" is meant release of the active component (e.g perfume) over a longer period of time than by the use of the active (e.g., perfume) itself.

Still in another aspect of the invention, there is provided the use of the product of the invention for the manufacture of a laundry and/or cleaning composition for delivering residual fragrance and fabric care, in particular color care, onto the fabrics on which it is applied.

5

The following are synthesis examples of compounds as defined in the present invention:

I-Synthesis of ethyl 4-aminobenzoate with 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde

10

To an ice cooled stirred solution of 10 g of 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde (0.07 mol) in 35 mL EtOH and molecular sieves (4Å, 20 g) 1eq of the amine was added via an addition funnel. The reaction mixture was stirred under nitrogen atmosphere and protected from light. After 6 days the mixture was filtrated and the solvent was removed. The yield of imine formation is about 90%.

15

Similar results were obtained where the 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde was replaced by bourgeonal, trans-2-nonenal, or trans-2-hexenal.

20

II-Synthesis of 1,4-bis-(3-aminopropyl)-piperazine with α -Damascone

In order to substitute both primary amine groups with a perfume, 2eq of perfume were used for 1eq of amino functional polymer. To an ice cooled stirred solution of 1mmol of α -Damascone in 6 mL EtOH and molecular sieves (4Å, 20 g), 0.5eq of 1,4-bis-(3-aminopropyl)-piperazine was added via an addition funnel. The reaction mixture was stirred under nitrogen atmosphere and protected from light. After the disappearance of the absorption peak from the NMR spectrum of the free perfume raw material (from 3 to 16 hours), the mixture was filtrated and the solvent was removed by vacuum distillation. The yield of β -aminoketone formation is about 90%.

25

30

Similar results were obtained where the α -Damascone was replaced by 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde, vertocitral, bourgeonal, δ -Damascone or citronellal. In these instances, Schiff-bases are formed.

35

III-Synthesis of Lupasol with Damascones and 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde

The β -amino ketone from Lupasol G100 (commercially available from BASF with a content of 50 % water, 50 % Lupasol G100 (Mw. 5000)) and α -Damascone (or δ -Damascone) was prepared using any one of these three different procedures described as follows:

1. Commercially available Lupasol G100 was dried using the following procedure: 20 g of the Lupasol solution was dried at the rotating evaporator during several hours. The obtained residue, still containing about 4.5 g of H_2O , was azeotropically distilled at the rotating evaporator using toluene. The residue was then placed in the desiccator dried at 60 °C (using P_2O_5 as water absorbing material). On basis of the obtained weight we concluded that the oil contained less then 10 % H_2O . On basis of the NMR-spectra we concluded that this is probably less then 5 %. This dried sample was then used in the preparation of β -amino ketones.

1.38 g of the dried Lupasol G100 obtained above was dissolved in 7 ml. ethanol. The solution was stirred gently with a magnetic stirrer during a few minutes before 2 g Na_2SO_4 (anhydrous) was added. After stirring again for a few minutes 2.21 g α -Damascone was added over a period of 1 minute. After two days reaction, the mixture was filtrated over a Celite filter (vide supra), and the residue washed thoroughly with ethanol. About 180 ml. of a light foaming filtrate was obtained. This was concentrated until dryness using a rotating evaporator and dried over P_2O_5 in an desiccator at room temperature. About 3.5 of a colorless oil was obtained.

2. 4.3 g Lupasol G100 solution was without drying dissolved in 10 ml. ethanol. The solution was stirred with a magnetic stirrer during a few minutes before 3.47 g α -Damascone was added over a 1.5 minutes period. After two days reaction at room temperature the reaction mixture was filtrated over Celite (vide supra) and the residue washed thoroughly with ethanol. The filtrate (200 ml., light foaming) was concentrated at the evaporator and dried in an desiccator (P_2O_5 as drying agent) at room temperature. About 5.9 g of a colorless oil was obtained.

3. To 3.0 g of Lupasol G100 solution (used as such) was added 2.41 g α -Damascone. The mixture was stirred without using solvent. After stirring for 4 days the obtained oil was dissolved in 100 ml. THF, dried with MgSO_4 , filtrated and the filtrate concentrated at the rotating evaporator. After drying in the desiccator (P_2O_5) at room temperature about 4 g of a colorless oil was obtained. This oil still contained about 13 % (w/w) of THF, even after a prolonged drying (3 days).

The product obtained from the three procedures had identical NMR-spectra.

Similar results are obtained where Lupasol G35 or Lupasol HF is used instead of Lupasol G100.

Similar results were obtained where the α -Damascone was replaced by 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde.

Still another possible route of synthesis is by using Lupasol P. The β -amino ketone from Lupasol P and α -Damascone was prepared using the procedure described as follows:

1.8g Lupasol P solution (50 % H_2O , 50 % Lupasol Mw. 750000, as obtained from BASF) was dissolved in 7 ml ethanol, the solution was stirred for a few minutes with a magnetic stirrer before 1.44 g α -Damascone was added. After three days the reaction mixture was filtrated over a celite filter (vide supra) and the residue washed thoroughly with ethanol. After concentrating of the filtrate and drying of the obtained oil in the desiccator (P_2O_5) at room temperature, about 3 g of the reaction product between Lupasol and α -Damascone was obtained.

Processing method

1)-Processing method for spray-on

Processing of the amine reaction product with the carrier is done as hereinbefore described. In particular, 20g of amine reaction product as above synthesised is mixed with 10 to 80g of a carrier, e.g. hydrophobic perfume composition, dipropylene glycol, diethylphthalate in an Ultra Turrax until an homogenous mixture is obtained. The temperature of mixing is about room temperature, and

the speed of the mixer is sufficient to maintain such temperature substantially constant. Temperature and time will depend on the nature of the carrier but are a conventional step to skilled man. Of course, the amount of carrier relative to the amine reaction product will depend on the desired sprayability and the end viscosity of the resulting mixture. In the examples of the present invention, the viscosity was below 500Cps, as measured on a rheometer, TA Instrument CSL²₁₀₀ at a temperature of 25C with a gap setting of 500 microns.

2)-Processing method for coating

Processing of the amine reaction product with the carrier is done as hereinbefore described. In particular, 20g of amine reaction product as above synthesised is mixed in an Ultra Turrax containing 20g of carrier, e.g. methanol or perfume mixture or composition for 5 minutes, the temperature of mixing being to about room temperature, and the speed of the mixer being sufficient so as to maintain such temperature substantially constant. Temperature and time will depend on the nature of the carrier but are a conventional step to skilled man.

When use of a perfume mixture is made, the composition in % by weight of a typical hydrophobic perfume mixture is as follows:

| | |
|--|-----|
| Citronellol | 7 |
| Geraniol | 7 |
| Linalool | 7 |
| Para Tertiary Butyl Cyclohexyl Acetate | 10 |
| Phenyl Ethyl Alcohol | 19 |
| Habanolide | 4.5 |
| Para Methoxy Acetophenone | 1.5 |
| Benzyl Acetate | 4 |
| Eugenol | 2 |
| Phenyl Ethyl Acetate | 5 |
| Verdyl Acetate | 6 |
| Verdyl Propionate | 4 |
| Hexyl Cinnamic Aldehyde | 3 |
| Ionone Gamma Methyl | 2 |
| Methyl Cedrylone | 10 |
| P.T. Bucinal | 7 |

| | |
|--------------------------|---|
| Para Cresyl Methyl Ether | 1 |
|--------------------------|---|

When desired, the obtained processed amine reaction product from above is subsequently coated. In particular, the resulting mixture is maintained at a temperature substantially equal to the melting point of the carrier material and poured onto the coating material and agglomerated in an electrical mixer like a Braun Mixer. Care is also taken that the temperature during the mixing does not substantially exceed the melting point of the carrier material. For example, 100g of a mixture containing the methanol or perfume mixture and 20g of the amine reaction product is poured at room temperature into a Braun Mixer containing 300g of carbonate. The mixing of the ingredients is carried out for about 5 minutes. Again, temperature and time will depend on the nature of the coating agent but are conventional step to skilled man.

Once the coated particle is obtained, it may be desired to remove the methanol by vacuum distillation.

Abbreviations used in the following laundry and cleaning composition Examples

In the laundry and cleaning compositions, the abbreviated component identifications have the following meanings:

| | | |
|--------------|---|--|
| DEQA | : | Di-(tallowyl-oxy-ethyl) dimethyl ammonium chloride |
| DTDMAC | : | Ditallow dimethylammonium chloride |
| DEQA (2) | : | Di-(soft-tallowyloxyethyl) hydroxyethyl methyl ammonium methylsulfate. |
| DTDMAMS | : | Ditallow dimethyl ammonium methylsulfate. |
| SDASA | : | 1:2 ratio of stearyl dimethyl amine:triple-pressed stearic acid. |
| Fatty acid | : | Stearic acid of IV=0 |
| Electrolyte | : | Calcium chloride |
| PEG | : | Polyethylene Glycol 4000 |
| Neodol 45-13 | : | C14-C15 linear primary alcohol ethoxylate, sold by Shell Chemical CO. |

| | | | |
|----|-------------------|---|--|
| | Silicone antifoam | : | Polydimethylsiloxane foam controller with siloxane-oxyalkylene copolymer as dispersing agent with a ratio of said foam controller to said dispersing agent of 10:1 to 100:1. |
| | PEI | : | Polyethyleneimine with an average molecular weight of 1800 and an average ethoxylation degree of 7 ethyleneoxy residues per nitrogen |
| 5 | HEDP | : | 1,1-hydroxyethane diphosphonic acid |
| | LAS | : | Sodium linear C ₁₁₋₁₃ alkyl benzene sulfonate |
| | TAS | : | Sodium tallow alkyl sulfate |
| | CxyAS | : | Sodium C _{1x} - C _{1y} alkyl sulfate |
| 10 | C46SAS | : | Sodium C ₁₄ - C ₁₆ secondary (2,3) alkyl sulfate |
| | CxyEzS | : | Sodium C _{1x} -C _{1y} alkyl sulfate condensed with z moles of ethylene oxide |
| | CxyEz | : | C _{1x} -C _{1y} predominantly linear primary alcohol condensed with an average of z moles of ethylene oxide |
| 15 | QAS | : | R ₂ .N ⁺ (CH ₃) ₂ (C ₂ H ₄ OH) with R ₂ = C ₁₂ - C ₁₄ |
| | QAS 1 | : | R ₂ .N ⁺ (CH ₃) ₂ (C ₂ H ₄ OH) with R ₂ = C ₈ - C ₁₁ |
| | APA | : | C ₈ - C ₁₀ amido propyl dimethyl amine |
| | Soap | : | Sodium linear alkyl carboxylate derived from an |
| 20 | | | 80/20 mixture of tallow and coconut fatty acids |
| | STS | : | Sodium toluene sulphonate |
| | CFAA | : | C ₁₂ -C ₁₄ (coco) alkyl N-methyl glucamide |
| | TFAA | : | C ₁₆ -C ₁₈ alkyl N-methyl glucamide |
| | TPKFA | : | C ₁₂ -C ₁₄ topped whole cut fatty acids |
| 25 | STPP | : | Anhydrous sodium tripolyphosphate |
| | TSPP | : | Tetrasodium pyrophosphate |
| | Zeolite A | : | Hydrated sodium aluminosilicate of formula Na ₁₂ (AlO ₂ SiO ₂) ₁₂ .27H ₂ O having a primary particle size in the range from 0.1 to 10 micrometers (weight expressed on an anhydrous basis) |
| 30 | NaSKS-6 | : | Crystalline layered silicate of formula δ- Na ₂ Si ₂ O ₅ |
| | Citric acid | : | Anhydrous citric acid |

| | | | |
|----|-----------------|---|--|
| | Borate | : | Sodium borate |
| | Carbonate | : | Anhydrous sodium carbonate with a particle size between 200µm and 900µm |
| 5 | Bicarbonate | : | Anhydrous sodium bicarbonate with a particle size distribution between 400µm and 1200µm |
| | Silicate | : | Amorphous sodium silicate ($\text{SiO}_2:\text{Na}_2\text{O} = 2.0:1$) |
| | Sulfate | : | Anhydrous sodium sulfate |
| | Mg sulfate | : | Anhydrous magnesium sulfate |
| 10 | Citrate | : | Tri-sodium citrate dihydrate of activity 86.4% with a particle size distribution between 425µm and 850µm |
| | MA/AA | : | Copolymer of 1:4 maleic/acrylic acid, average molecular weight about 70,000 |
| | MA/AA (1) | : | Copolymer of 4:6 maleic/acrylic acid, average molecular weight about 10,000 |
| 15 | AA | : | Sodium polyacrylate polymer of average molecular weight 4,500 |
| | CMC | : | Sodium carboxymethyl cellulose |
| 20 | Cellulose ether | : | Methyl cellulose ether with a degree of polymerization of 650 available from Shin Etsu Chemicals |
| | Protease | : | Proteolytic enzyme, having 3.3% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Savinase |
| 25 | Protease I | : | Proteolytic enzyme, having 4% by weight of active enzyme, as described in WO 95/10591, sold by Genencor Int. Inc. |
| | Alcalase | : | Proteolytic enzyme, having 5.3% by weight of active enzyme, sold by NOVO Industries A/S |
| 30 | Cellulase | : | Cellulytic enzyme, having 0.23% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Carezyme |
| | Amylase | : | Amylolytic enzyme, having 1.6% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Termamyl 120T |

| | | | |
|----|---------------------------|---|--|
| | Lipase | : | Lipolytic enzyme, having 2.0% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Lipolase |
| 5 | Lipase (1) | : | Lipolytic enzyme, having 2.0% by weight of active enzyme, sold by NOVO Industries A/S under the tradename Lipolase Ultra |
| | Endolase | : | Endoglucanase enzyme, having 1.5% by weight of active enzyme, sold by NOVO Industries A/S |
| 10 | PB4 | : | Sodium perborate tetrahydrate of nominal formula $\text{NaBO}_2 \cdot 3\text{H}_2\text{O} \cdot \text{H}_2\text{O}_2$ |
| | PB1 | : | Anhydrous sodium perborate bleach of nominal formula $\text{NaBO}_2 \cdot \text{H}_2\text{O}_2$ |
| | Percarbonate | : | Sodium percarbonate of nominal formula $2\text{Na}_2\text{CO}_3 \cdot 3\text{H}_2\text{O}_2$ |
| 15 | NOBS | : | Nonanoyloxybenzene sulfonate in the form of the sodium salt |
| | NAC-OBS | : | (6-nonamidocaproyl) oxybenzene sulfonate |
| | TAED | : | Tetraacetythylenediamine |
| | DTPA | : | Diethylene triamine pentaacetic acid |
| 20 | DTPMP | : | Diethylene triamine penta (methylene phosphonate), marketed by Monsanto under the Tradename Dequest 2060 |
| | EDDS | : | Ethylenediamine-N,N'-disuccinic acid, (S,S) isomer in the form of its sodium salt. |
| 25 | Photoactivated bleach (1) | : | Sulfonated zinc phthlocyanine encapsulated in dextrin soluble polymer |
| | Photoactivated bleach (2) | : | Sulfonated alumino phthlocyanine encapsulated in dextrin soluble polymer |
| | Brightener 1 | : | Disodium 4,4'-bis(2-sulphostyryl)biphenyl |
| 30 | Brightener 2 | : | Disodium 4,4'-bis(4-anilino-6-morpholino-1,3,5-triazin-2-yl)amino) stilbene-2,2'-disulfonate |
| | HEDP | : | 1,1-hydroxyethane diphosphonic acid |
| | PEGx | : | Polyethylene glycol, with a molecular weight of x (typically 4,000) |
| 35 | PEO | : | Polyethylene oxide, with an average molecular weight of 50,000 |

| | | | |
|----|-------------------|---|---|
| | TEPAE | : | Tetraethylenepentaamine ethoxylate |
| | PVI | : | Polyvinyl imidosole, with an average molecular weight of 20,000 |
| 5 | PVP | : | Polyvinylpyrrolidone polymer, with an average molecular weight of 60,000 |
| | PVNO | : | Polyvinylpyridine N-oxide polymer, with an average molecular weight of 50,000 |
| | PVPVI | : | Copolymer of polyvinylpyrrolidone and vinylimidazole, with an average molecular weight of 20,000 |
| 10 | QEA | : | bis((C ₂ H ₅ O)(C ₂ H ₄ O) _n)(CH ₃) -N ⁺ -C ₆ H ₁₂ -N ⁺ -(CH ₃) bis((C ₂ H ₅ O)-(C ₂ H ₄ O)) _n , wherein n = from 20 to 30 |
| | SRP 1 | : | Anionically end capped poly esters |
| | SRP 2 | : | Diethoxylated poly (1, 2 propylene terephthalate) short block polymer |
| 15 | PEI | : | Polyethyleneimine with an average molecular weight of 1800 and an average ethoxylation degree of 7 ethyleneoxy residues per nitrogen |
| | Silicone antifoam | : | Polydimethylsiloxane foam controller with siloxane-oxyalkylene copolymer as dispersing agent with a ratio of said foam controller to said dispersing agent of 10:1 to 100:1 |
| 20 | Opacifier | : | Water based monostyrene latex mixture, sold by BASF Aktiengesellschaft under the tradename Lytron 621 |
| 25 | Wax | : | Paraffin wax |
| | PA30 | : | Polyacrylic acid of average molecular weight of between about 4,500 - 8,000. |
| | 480N | : | Random copolymer of 7:3 acrylate/methacrylate, average molecular weight about 3,500. |
| | Polygel/carbopol | : | High molecular weight crosslinked polyacrylates. |
| | Metasilicate | : | Sodium metasilicate (SiO ₂ :Na ₂ O ratio = 1.0). |
| | Nonionic | : | C ₁₃ -C ₁₅ mixed ethoxylated/propoxylated fatty alcohol with an average degree of ethoxylation of 3.8 and an average degree of propoxylation of 4.5. |
| | Neodol 45-13 | : | C ₁₄ -C ₁₅ linear primary alcohol ethoxylate, sold by Shell Chemical CO. |

| | | |
|----------|---|--|
| MnTACN | : | Manganese 1,4,7-trimethyl-1,4,7-triazacyclononane. |
| PAAC | : | Pentaamine acetate cobalt(III) salt. |
| Paraffin | : | Paraffin oil sold under the tradename Winog 70 by Wintershall. |
| NaBz | : | Sodium benzoate. |
| BzP | : | Benzoyl Peroxide. |
| SCS | : | Sodium cumene sulphonate. |
| BTA | : | Benzotriazole. |
| PH | : | Measured as a 1% solution in distilled water at 20°C. |
| PARP1 | : | Processed amine reaction product of ethyl 4-aminobenzoate with 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde as made from Synthesis example I, mixed with a methanol carrier, optionally agglomerated with a coating agent as specified hereinafter in the examples, and followed by removal of the methanol, according to processing method above described. |
| PARP2 | : | Processed amine reaction product of Lupasol G35 with α -Damascone as made from Synthesis example III, mixed with a perfume mixture carrier as described herein before and sprayed on finished product, according to processing method above described. |
| PARP3 | : | Processed amine reaction product of Lupasol HF with δ -Damascone as made from Synthesis example III, mixed with a methanol carrier, optionally agglomerated with a coating agent as specified hereinafter in the examples, and followed by removal of the methanol, according to processing method above described. |
| PARP4 | : | Processed amine reaction product of LupasolG100 with 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde as made from Synthesis example III, mixed with a perfume mixture carrier as described herein before and sprayed on finished product, according to processing method above described. |
| Clay I | : | Bentonite clay |
| Clay II | : | Smectite clay |

- Flocculating agent I: polyethylene oxide of average molecular weight of between 200,000 and 400,000
- Flocculating agent II: polyethylene oxide of average molecular weight of between 400,000 and 1,000,000
- 5 Flocculating agent III: polymer of acrylamide and/ or acrylic acid of average molecular weight of 200,000 and 400,000
- DOBS: Decanoyl oxybenzene sulfonate in the form of the sodium salt
- SRP 3 : Polysaccharide soil release polymer
- 10 SRP 4 : Nonionically end capped poly esters
- Polymer : Polyvinylpyrrolidone K90 available from BASF under the tradename Luviskol K90
- 15 Dye fixative : Dye fixative commercially available from Clariant under the tradename Cartafix CB
- Polyamine : 1,4-Bis-(3-aminopropyl)piperazine
- Bayhibit AM : 2-Phosphonobutane-1,2,4-tricarboxylic acid
- 20 commercially available from Bayer
- Fabric softener active: Di-(canoloyl-oxy-ethyl)hydroxyethyl methyl ammonium methylsulfate
- HPBDC : Hydroxypropyl beta-cyclodextrin
- RAMEB : Randomly methylated beta-cyclodextrin
- 25 Bardac 2050 : Dioctyl dimethyl ammonium chloride, 50% solution
- Bardac 22250 : Didecyl dimethyl ammonium chloride, 50% solution
- Genamin C100 : Coco fatty amine ethoxylated with 10 moles ethylene oxide and commercially available from Clariant
- 30 Genapol V4463 : Coco alcohol ethoxylated with 10 moles ethylene oxide and commercially available from Clariant
- Silwet 7604: Polyalkyleneoxide polysiloxanes of MW 4000 of formula
- 35 $R-(CH_3)_2SiO-[(CH_3)_2SiO]_a-[(CH_3)(R)SiO]_b-Si(CH_3)_2-R$, wherein average $a+b$ is 21, and commercially available from Osi Specialties, Inc., Danbury, Connecticut

Silwet 7600: Polyalkyleneoxide polysiloxanes of MW 4000, of formula
 $R-(CH_3)_2SiO-[(CH_3)_2SiO]_a-[(CH_3)(R)SiO]_b-Si(CH_3)_2-R$,
 wherein average $a+b$ is 11, and commercially available from Osi Specialties, Inc., Danbury, Connecticut

In the following formulation examples all levels are quoted as % by weight of the composition unless otherwise stated, and incorporation of the processed amine reaction product so called herein after "PARP" in the fully formulated composition is carried out by either dry addition in the composition or spray-on as defined herein before depending on its end form. The term in bracket for the PARP in the formulation examples refers, when present, to the type of coating (c) for carbonate coating and (s) for starch coating.

Example 1

The following high density granular laundry detergent compositions are in accord with the invention:

| | A | B | C | D | E |
|-------------|------|------|------|------|------|
| LAS | 6.0 | 6.0 | 8.0 | 8.0 | 8.0 |
| TAS | 1.0 | 0.1 | - | 0.5 | - |
| C46(S)AS | - | - | 2.0 | 2.5 | - |
| C25AS | 4.5 | 5.5 | - | - | - |
| C68AS | - | - | 2.0 | 5.0 | 7.0 |
| C25E5 | 4.6 | 4.6 | - | - | 3.4 |
| C25E7 | - | - | 3.4 | 3.4 | 1.0 |
| C25E3S | 5.0 | 4.5 | - | - | - |
| QAS | - | - | - | 0.8 | - |
| QAS (I) | 0.5 | 1.0 | - | - | - |
| Zeolite A | 20.0 | 18.1 | 18.1 | 18.0 | 14.1 |
| Citric acid | - | 2.5 | - | - | - |
| Carbonate | 10.0 | 13.0 | 13.0 | 13.0 | 25.0 |
| SKS-6 | - | 10.0 | - | - | - |
| Silicate | 0.5 | 0.3 | 1.4 | 1.4 | 3.0 |
| Citrate | - | - | - | 1.0 | - |

| | | | | | |
|--------------------------------|-----------|-----------|--------|--------|--------|
| Sulfate | - | - | 26.1 | 26.1 | 26.1 |
| Mg sulfate | - | 0.2 | 0.3 | - | - |
| MA/AA | 1.0 | 1.0 | 0.3 | 0.3 | 0.3 |
| CMC | 0.4 | 0.4 | 0.2 | 0.2 | 0.2 |
| PB4 | - | - | 9.0 | 9.0 | 5.0 |
| Percarbonate | 18.0 | 18.0 | - | - | - |
| TAED | 3.9 | 4.2 | 1.5 | 0.4 | 1.5 |
| NAC-OBS | - | - | - | 2.0 | 1.0 |
| DTPMP | - | - | 0.25 | 0.25 | 0.25 |
| SRP 2 | - | 0.2 | - | - | - |
| EDDS | 0.5 | 0.5 | - | 0.25 | 0.4 |
| CFAA | - | - | - | 1.0 | - |
| HEDP | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 |
| QEA | - | 0.5 | - | - | - |
| Protease I | - | - | - | - | 0.26 |
| Protease | 1.5 | 1.0 | 0.26 | 0.26 | - |
| Cellulase | 0.3 | 0.3 | 0.3 | - | - |
| Amylase | 0.5 | 0.5 | 0.1 | 0.1 | 0.1 |
| Lipase (1) | 0.5 | 0.5 | 0.3 | - | - |
| Photoactivated bleach (ppm) | 20 ppm | 20 ppm | 15 ppm | 15 ppm | 15 ppm |
| Brightener 1 | 0.09 | 0.09 | 0.09 | 0.09 | 0.09 |
| Perfume spray on | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 |
| PARP 1 | 0.9(c) | 1.2(c) | 1.0(c) | 0.4(s) | - |
| PARP 3 | 0.2 | 0.3 | - | - | 1.0 |
| Silicone antifoam | 0.3 | 0.3 | 0.5 | 0.5 | 0.5 |
| Misc/minors to 100% | | | | | |
| Density in g/litre | 850 | 850 | 850 | 850 | 850 |

| | F | G | H | I |
|--------------|------|------|------|------|
| LAS | 2.0 | 6.0 | 6.0 | 5.0 |
| TAS | 0.5 | 1.0 | 0.1 | 1.5 |
| C25AS | 7.0 | 4.5 | 5.5 | 2.5 |
| C68AS | - | - | - | 0.2 |
| C25E5 | 10.0 | 4.6 | 4.6 | 2.6 |
| C25E3S | 2.0 | 5.0 | 4.5 | 0.5 |
| QAS (I) | 0.8 | 0.5 | 1.0 | 1.5 |
| Zeolite A | 18.1 | 20.0 | 18.1 | 16.2 |
| Citric acid | 2.5 | - | 2.5 | 1.5 |
| Carbonate | 10.0 | 10.0 | 13.0 | 20.6 |
| SKS-6 | 10.0 | - | 10.0 | 4.3 |
| Silicate | 0.3 | 0.5 | 0.3 | - |
| Citrate | 3.0 | - | - | 1.4 |
| Sulfate | 6.0 | - | - | - |
| Mg sulfate | 0.2 | - | 0.2 | 0.03 |
| MA/AA | 4.0 | 1.0 | 1.0 | 0.6 |
| CMC | 0.2 | 0.4 | 0.4 | 0.3 |
| Percarbonate | - | 18.0 | 18.0 | 9.0 |
| TAED | - | 3.9 | 4.2 | 3.2 |
| DTPMP | 0.25 | - | - | - |
| SRP 2 | 0.2 | - | 0.2 | - |
| EDDS | - | 0.5 | 0.5 | 0.1 |
| CFAA | 2.0 | - | - | - |
| TFAA | - | - | - | 1.1 |
| HEDP | 0.3 | 0.4 | 0.4 | 0.3 |
| QEA | 0.2 | - | 0.5 | - |
| Protease I | 1.0 | - | - | 0.3 |
| Protease | - | 1.5 | 1.0 | - |
| Cellulase | 0.3 | 0.3 | 0.3 | 0.3 |
| Amylase | 0.4 | 0.5 | 0.5 | 0.1 |
| Lipase (1) | 0.5 | 0.5 | 0.5 | 0.1 |

| | | | | |
|-----------------------------|--------|---------|--------|--------|
| Photoactivated bleach (ppm) | - | 20 ppm | 20 ppm | 20 ppm |
| PVNO/PVPVI | 0.1 | - | - | - |
| Brightener 1 | - | 0.09 | 0.09 | 0.01 |
| Brightener 2 | - | - | - | 0.09 |
| Perfume spray on | 0.4 | 0.4 | 0.4 | 0.4 |
| PARP 1 | - | 0.5(c) | 1.0(c) | - |
| PARP 3 | 0.8(c) | 0.4 (s) | - | 0.4(c) |
| Silicone antifoam | - | 0.3 | 0.3 | 0.3 |
| Clay II | - | - | - | 12.0 |
| Flocculating agent I | - | - | - | 0.3 |
| Glycerol | - | - | - | 0.6 |
| Wax | - | - | - | 0.4 |
| Misc/minors to 100% | | | | |
| Density in g/litre | 850 | 850 | 850 | 850 |

Example 2

The following granular laundry detergent compositions of particular utility under European machine wash conditions are in accord with the invention:

5

| | A | B | C | D | E | F |
|-------------|------|------|------|------|------|------|
| LAS | 5.5 | 7.5 | 5.0 | 5.0 | 6.0 | 7.0 |
| TAS | 1.25 | 1.86 | - | 0.8 | 0.4 | 0.3 |
| C24AS/C25AS | - | 2.24 | 5.0 | 5.0 | 5.0 | 2.2 |
| C25E3S | - | 0.76 | 1.0 | 1.5 | 3.0 | 1.0 |
| C45E7 | 3.25 | - | - | - | - | 3.0 |
| TFAA | - | - | 2.0 | - | - | - |
| C25E5 | - | 5.5 | - | - | - | - |
| QAS | 0.8 | - | - | - | - | - |
| QAS II | - | 0.7 | 1.0 | 0.5 | 1.0 | 0.7 |
| STPP | 19.7 | - | - | - | - | - |
| Zeolite A | - | 19.5 | 25.0 | 19.5 | 20.0 | 17.0 |

| | | | | | | |
|---------------------------------|---------|--------|-----|------|--------|--------|
| NaSKS-6/citric acid (79:21) | - | 10.6 | - | 10.6 | - | - |
| NaSKS-6 | - | - | 9.0 | - | 10.0 | 10.0 |
| Carbonate | 6.1 | 10.0 | 9.0 | 10.0 | 10.0 | 18.0 |
| Bicarbonate | - | 2.0 | 7.0 | 5.0 | - | 2.0 |
| Silicate | 6.8 | - | - | 0.3 | 0.5 | - |
| Citrate | - | - | 4.0 | 4.0 | - | - |
| Sulfate | 39.8 | - | - | 5.0 | - | 12.0 |
| Mg sulfate | - | - | 0.1 | 0.2 | 0.2 | - |
| MA/AA | 0.5 | 1.6 | 3.0 | 4.0 | 1.0 | 1.0 |
| CMC | 0.2 | 0.4 | 1.0 | 1.0 | 0.4 | 0.4 |
| PB4 | 5.0 | 12.7 | - | - | - | - |
| Percarbonate | - | - | - | - | 18.0 | 15.0 |
| TAED | 0.5 | 3.1 | - | - | 5.0 | - |
| NAC-OBS | 1.0 | 3.5 | - | - | - | 2.5 |
| DTPMP | 0.25 | 0.2 | 0.3 | 0.4 | - | 0.2 |
| HEDP | - | 0.3 | - | 0.3 | 0.3 | 0.3 |
| QEA | - | - | 1.0 | 1.0 | 1.0 | - |
| Protease I | - | - | - | 0.5 | 1.2 | - |
| Protease | 0.26 | 0.85 | 0.9 | 1.0 | - | 0.7 |
| Lipase (1) | 0.15 | 0.15 | 0.3 | 0.3 | 0.3 | 0.2 |
| Cellulase | 0.28 | 0.28 | 0.2 | 0.2 | 0.3 | 0.3 |
| Amylase | 0.1 | 0.1 | 0.4 | 0.4 | 0.6 | 0.2 |
| PVNO/PVPVI | - | - | 0.2 | 0.2 | - | - |
| PVP | 0.9 | 1.3 | - | - | - | 0.9 |
| SRP 1 | - | - | 0.2 | 0.2 | 0.2 | - |
| Photoactivated bleach (1) (ppm) | 15 ppm | 27 ppm | - | - | 20 ppm | 20 ppm |
| Photoactivated bleach (2) (ppm) | 15 ppm | - | - | - | - | - |
| Brightener 1 | 0.08 | 0.19 | - | - | 0.09 | 0.15 |
| Brightener 2 | - | 0.04 | - | - | - | - |
| Perfume | 0.3 | 0.3 | 0.4 | 0.3 | 0.4 | 0.3 |
| PARP1 | 1.0 (c) | 2.0(c) | 1.5 | - | - | - |
| PARP2 | - | - | 0.2 | 0.1 | 0.8 | 0.4 |

| | | | | | | |
|---------------------|-----|-----|-----|-----|-----|-----|
| Silicone antifoam | 0.5 | 2.4 | 0.3 | 0.5 | 0.3 | 2.0 |
| Minors/misc to 100% | | | | | | |
| Density in g/litre | 750 | 750 | 750 | 750 | 750 | 750 |

Example 3

The following detergent formulations of particular utility under European machine wash conditions were prepared in accord with the invention.

5

| | A | B | C | D |
|-------------------|------|------|------|------|
| Blown powder | | | | |
| LAS | 6.0 | 5.0 | 11.0 | 6.0 |
| TAS | 2.0 | - | - | 2.0 |
| Zeolite A | 24.0 | - | - | 20.0 |
| STPP | - | 27.0 | 24.0 | - |
| Sulfate | 4.0 | 6.0 | 13.0 | - |
| MA/AA | 1.0 | 4.0 | 6.0 | 2.0 |
| Silicate | 1.0 | 7.0 | 3.0 | 3.0 |
| CMC | 1.0 | 1.0 | 0.5 | 0.6 |
| Brightener 1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Silicone antifoam | 1.0 | 1.0 | 1.0 | 0.3 |
| DTPMP | 0.4 | 0.4 | 0.2 | 0.4 |
| Spray on | | | | |
| Brightener | 0.02 | - | - | 0.02 |
| C45E7 | - | - | - | 5.0 |
| C45E2 | 2.5 | 2.5 | 2.0 | - |
| C45E3 | 2.6 | 2.5 | 2.0 | - |
| Perfume | 0.5 | 0.3 | 0.5 | 0.2 |
| Silicone antifoam | 0.3 | 0.3 | 0.3 | - |
| Dry additives | | | | |
| QEA | - | - | - | 1.0 |
| EDDS | 0.3 | - | - | - |
| Sulfate | 2.0 | 3.0 | 5.0 | 10.0 |
| Carbonate | 6.0 | 13.0 | 11.0 | 14.0 |
| Citric acid | 2.5 | - | - | 2.0 |

| | | | | |
|--------------------|---------|--------|--------|------|
| QAS II | 0.5 | - | - | 0.5 |
| SKS-6 | 10.0 | - | - | - |
| Percarbonate | 18.5 | - | - | - |
| PB4 | - | 18.0 | 10.0 | 21.5 |
| TAED | 2.0 | 2.0 | - | 2.0 |
| NAC-OBS | 3.0 | 2.0 | 4.0 | - |
| Protease | 1.0 | 1.0 | 1.0 | 1.0 |
| Lipase | - | 0.4 | - | 0.2 |
| Lipase (1) | 0.4 | - | 0.4 | - |
| Amylase | 0.2 | 0.2 | 0.2 | 0.4 |
| Brightener 1 | 0.05 | - | - | 0.05 |
| PARP3 | 0.3 (c) | 1.0(c) | 0.4(s) | 1.2 |
| Misc/minor to 100% | | | | |

Example 4

The following granular detergent formulations were prepared in accord with the invention.

5

| | A | B | C | D | E | F |
|--------------|------|------|------|------|------|------|
| Blown powder | | | | | | |
| LAS | 23.0 | 8.0 | 7.0 | 9.0 | 7.0 | 7.0 |
| TAS | - | - | - | - | 1.0 | - |
| C45AS | 6.0 | 6.0 | 5.0 | 8.0 | - | - |
| C45AES | - | 1.0 | 1.0 | 1.0 | - | - |
| C45E35 | - | - | - | - | 2.0 | 4.0 |
| Zeolite A | 10.0 | 18.0 | 14.0 | 12.0 | 10.0 | 10.0 |
| MA/AA | - | 0.5 | - | - | - | 2.0 |
| MA/AA (1) | 7.0 | - | - | - | - | - |
| AA | - | 3.0 | 3.0 | 2.0 | 3.0 | 3.0 |
| Sulfate | 5.0 | 6.3 | 14.3 | 11.0 | 15.0 | 19.3 |
| Silicate | 10.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 |
| Carbonate | 13.0 | 19.0 | 8.0 | 20.7 | 8.0 | 6.0 |
| PEG 4000 | 0.4 | 1.5 | 1.5 | 1.0 | 1.0 | 1.0 |
| DTPA | - | 0.9 | 0.5 | - | - | 0.5 |

| | | | | | | |
|----------------------|--------|--------|--------|--------|--------|--------|
| Brightener 2 | 0.3 | 0.2 | 0.3 | - | 0.1 | 0.3 |
| Spray on | | | | | | |
| C45E7 | - | 2.0 | - | - | 2.0 | 2.0 |
| C25E9 | 3.0 | - | - | - | - | - |
| C23E9 | - | - | 1.5 | 2.0 | - | 2.0 |
| Perfume | 0.3 | 0.3 | 0.3 | 2.0 | 0.3 | 0.3 |
| Agglomerates | | | | | | |
| C45AS | - | 5.0 | 5.0 | 2.0 | - | 5.0 |
| LAS | - | 2.0 | 2.0 | - | - | 2.0 |
| Zeolite A | - | 7.5 | 7.5 | 8.0 | - | 7.5 |
| Carbonate | - | 4.0 | 4.0 | 5.0 | - | 4.0 |
| PEG 4000 | - | 0.5 | 0.5 | - | - | 0.5 |
| Misc (water etc) | - | 2.0 | 2.0 | 2.0 | - | 2.0 |
| Dry additives | | | | | | |
| QAS (I) | - | - | - | - | 1.0 | - |
| Citric acid | - | - | - | - | 2.0 | - |
| PB4 | - | - | - | - | 12.0 | 1.0 |
| PB1 | 4.0 | 1.0 | 3.0 | 2.0 | - | - |
| Percarbonate | - | - | - | - | 2.0 | 10.0 |
| Carbonate | - | 5.3 | 1.8 | - | 4.0 | 4.0 |
| NOBS | 4.0 | - | 6.0 | - | - | 0.6 |
| Methyl cellulose | 0.2 | - | - | - | - | - |
| SKS-6 | 8.0 | - | - | - | - | - |
| STS | - | - | 2.0 | - | 1.0 | - |
| Cumene sulfonic acid | - | 1.0 | - | - | - | 2.0 |
| Lipase | 0.2 | - | 0.2 | - | 0.2 | 0.4 |
| Cellulase | 0.2 | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 |
| Amylase | 0.2 | - | 0.1 | - | 0.2 | - |
| Protease | 0.5 | 0.5 | 0.5 | 0.3 | 0.5 | 0.5 |
| PVPVI | - | - | - | - | 0.5 | 0.1 |
| PVP | - | - | - | - | 0.5 | - |
| PVNO | - | - | 0.5 | 0.3 | - | - |
| QEA | - | - | - | - | 1.0 | - |
| SRP1 | 0.2 | 0.5 | 0.3 | - | 0.2 | - |
| PARP3 | 0.2(c) | 0.3(c) | 1.0(c) | 0.5(c) | 1.0(c) | 0.1(s) |

| | | | | | | |
|---------------------|-----|-----|-----|-----|-----|---|
| Silicone antifoam | 0.2 | 0.4 | 0.2 | 0.4 | 0.1 | - |
| Mg sulfate | - | - | 0.2 | - | 0.2 | - |
| Misc/minors to 100% | | | | | | |

| | G | H | I | J |
|----------------------------|------|------|------|------|
| Blown powder | | | | |
| Clay I or II | 7.0 | 10.0 | 6.0 | 2.0 |
| LAS | 16.0 | 5.0 | 11.0 | 6.0 |
| TAS | - | 5.0 | - | 2.0 |
| Zeolite A | - | 20.0 | - | 10.0 |
| STPP | 24.0 | - | 14.0 | - |
| Sulfate | - | 2.0 | - | - |
| MA/AA | - | 2.0 | 1.0 | 1.0 |
| Silicate | 4.0 | 7.0 | 3.0 | - |
| CMC | 1.0 | - | 0.5 | 0.6 |
| Brightener 1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Carbonate | 10.0 | 10.0 | 20.0 | - |
| DTPMP | 0.4 | 0.4 | 0.2 | - |
| Spray on | | | | |
| Brightener 1 | 0.02 | - | - | 0.02 |
| C45E7 or E9 | - | - | 2.0 | 1.0 |
| C45E3 or E4 | - | - | 2.0 | 4.0 |
| Perfume | 0.5 | - | 0.5 | 0.2 |
| Silicone antifoam | 0.3 | - | - | - |
| PARP2 | 0.16 | 0.2 | 0.4 | 1.0 |
| Dry additives | | | | |
| Flocculating agent I or II | 0.3 | 1.0 | 1.0 | 0.5 |
| QEA | - | - | - | 1.0 |
| HEDP/ EDDS | 0.3 | - | - | - |
| Sulfate | 2.0 | - | - | - |
| Carbonate | 20.0 | 13.0 | 15.0 | 24.0 |
| Citric acid | 2.5 | - | - | 2.0 |
| QAS | - | - | 0.5 | 0.5 |
| NaSKS-6 | 3.5 | - | - | 5.0 |
| Percarbonate | - | - | - | 9.0 |

| | | | | |
|--------------------|------|-----|-----|------|
| PB4 | - | - | 5.0 | |
| NOBS | - | - | - | 1.3 |
| TAED | - | - | 2.0 | 1.5 |
| Protease | 1.0 | 1.0 | 1.0 | 1.0 |
| Lipase | - | 0.4 | - | 0.2 |
| Amylase | 0.2 | 0.2 | 0.2 | 0.4 |
| Brightener 2 | 0.05 | - | - | 0.05 |
| Perfume | 0.1 | 0.2 | 0.5 | - |
| Speckle | 1.2 | 0.5 | 2.0 | - |
| Misc/minor to 100% | | | | |

Example 5

The following nil bleach-containing detergent formulations of particular use in the washing of coloured clothing, according to the present invention were prepared:

5

| | A | B | C |
|---------------------|------|------|------|
| Blown Powder | | | |
| Zeolite A | 15.0 | 15.0 | - |
| Sulfate | 0.0 | 5.0 | - |
| LAS | 3.0 | 3.0 | - |
| DTPMP | 0.4 | 0.5 | - |
| CMC | 0.4 | 0.4 | - |
| MA/AA | 4.0 | 4.0 | - |
| Agglomerates | | | |
| C45AS | - | - | 11.0 |
| LAS | 6.0 | 5.0 | - |
| TAS | 3.0 | 2.0 | - |
| Silicate | 4.0 | 4.0 | - |
| Zeolite A | 10.0 | 15.0 | 13.0 |
| CMC | - | - | 0.5 |
| MA/AA | - | - | 2.0 |
| Carbonate | 9.0 | 7.0 | 7.0 |
| Spray On | | | |
| Perfume | 0.3 | 0.3 | 0.5 |
| C45E7 | 4.0 | 4.0 | 4.0 |

| | | | |
|---------------------|--------|--------|--------|
| C25E3 | 2.0 | 2.0 | 2.0 |
| PARP4 | 0.3 | 1.0 | 0.2 |
| Dry additives | | | |
| MA/AA | - | - | 3.0 |
| NaSKS-6 | - | - | 12.0 |
| Citrate | 10.0 | - | 8.0 |
| Bicarbonate | 7.0 | 3.0 | 5.0 |
| Carbonate | 6.0 | - | 7.0 |
| PVPVI/PVNO | 0.5 | 0.5 | 0.5 |
| Alcalase | 0.5 | 0.3 | 0.9 |
| Lipase | 0.4 | 0.4 | 0.4 |
| Amylase | 0.6 | 0.6 | 0.6 |
| Cellulase | 0.6 | 0.6 | 0.6 |
| PARP3 | 0.7(c) | 0.1(c) | 0.5(c) |
| Silicone antifoam | 5.0 | 5.0 | 5.0 |
| Dry additives | | | |
| Sulfate | 0.0 | 9.0 | 0.0 |
| Misc/minors to 100% | 100.0 | 100.0 | 100.0 |
| Density (g/litre) | 700 | 700 | 700 |

Example 6

The following granular detergent formulations were prepared in accord with the invention.

| | A | B | C | D |
|--------------|------|------|------|------|
| Base granule | | | | |
| Zeolite A | 30.0 | 22.0 | 24.0 | 10.0 |
| Sulfate | 10.0 | 5.0 | 10.0 | 7.0 |
| MA/AA | 3.0 | - | - | - |
| AA | - | 1.6 | 2.0 | - |
| MA/AA (1) | - | 12.0 | - | 6.0 |
| LAS | 14.0 | 10.0 | 9.0 | 20.0 |
| C45AS | 8.0 | 7.0 | 9.0 | 7.0 |
| C45AES | - | 1.0 | 1.0 | - |

| | | | | |
|--------------------|-----|--------|--------|--------|
| Silicate | - | 1.0 | 0.5 | 10.0 |
| Soap | - | 2.0 | - | - |
| Brightener 1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Carbonate | 6.0 | 9.0 | 10.0 | 10.0 |
| PEG 4000 | - | 1.0 | 1.5 | - |
| DTPA | - | 0.4 | - | - |
| Spray on | | | | |
| C25E9 | - | - | - | 5.0 |
| C45E7 | 1.0 | 1.0 | - | - |
| C23E9 | - | 1.0 | 2.5 | - |
| Perfume | - | 0.3 | 0.3 | - |
| PARP2 | 0.3 | 0.05 | 0.008 | - |
| Dry additives | | | | |
| Carbonate | 5.0 | 5.0 | 15.0 | 6.0 |
| PVPVI/PVNO | 0.5 | - | 0.3 | - |
| Protease | 1.0 | 1.0 | 1.0 | 0.5 |
| Lipase | 0.4 | - | - | 0.4 |
| Amylase | 0.1 | - | - | 0.1 |
| Cellulase | 0.1 | 0.2 | 0.2 | 0.1 |
| NOBS | - | 4.0 | - | 4.5 |
| PB1 | 1.0 | 5.0 | 1.5 | 6.0 |
| Sulfate | 4.0 | 5.0 | - | 5.0 |
| SRPI | - | 0.4 | - | - |
| PARP1 | - | 1.5(c) | 0.8(c) | - |
| PARP3 | - | 0.1 | - | 0.4(s) |
| Sud supressor | - | 0.5 | 0.5 | - |
| Misc/minor to 100% | | | | |

Example 7

The following granular detergent compositions were prepared in accord with the invention.

5

| | A | B | C |
|--------------|---|---|---|
| Blown powder | | | |

| | | | |
|-----------------------|--------|--------|--------|
| Zeolite A | 20.0 | - | 15.0 |
| STPP | - | 20.0 | - |
| Sulphate | - | - | 5.0 |
| Carbonate | - | - | 5.0 |
| TAS | - | - | 1.0 |
| LAS | 6.0 | 6.0 | 6.0 |
| C68AS | 2.0 | 2.0 | - |
| Silicate | 3.0 | 8.0 | - |
| MA/AA | 4.0 | 2.0 | 2.0 |
| CMC | 0.6 | 0.6 | 0.2 |
| Brightener 1 | 0.2 | 0.2 | 0.1 |
| DTPMP | 0.4 | 0.4 | 0.1 |
| STS | - | - | 1.0 |
| Spray on | | | |
| C45E7 | 5.0 | 5.0 | 4.0 |
| Silicone antifoam | 0.3 | 0.3 | 0.1 |
| Perfume | 0.2 | 0.2 | 0.3 |
| PARP4 | 1.2 | 1.0 | - |
| Dry additives | | | |
| QEA | - | - | 1.0 |
| Carbonate | 14.0 | 9.0 | 10.0 |
| PB1 | 1.5 | 2.0 | - |
| PB4 | 18.5 | 13.0 | 13.0 |
| TAED | 2.0 | 2.0 | 2.0 |
| QAS (I) | - | - | 1.0 |
| Photoactivated bleach | 15 ppm | 15 ppm | 15ppm |
| SKS-6 | - | - | 3.0 |
| Protease | 1.0 | 1.0 | 0.2 |
| Lipase | 0.2 | 0.2 | 0.2 |
| Amylase | 0.4 | 0.4 | 0.2 |
| Cellulase | 0.1 | 0.1 | 0.2 |
| Sulfate | 10.0 | 20.0 | 5.0 |
| PARP3 | 0.5(c) | 0.5(c) | 0.4(c) |
| Misc/minors to 100% | | | |
| Density (g/litre) | 700 | 700 | 700 |

Example 8

The following detergent compositions, according to the present invention were prepared:

5

| | A | B | C |
|----------------------|------|------|------|
| Blown Powder | | | |
| Zeolite A | 15.0 | 15.0 | 15.0 |
| Sulfate | 0.0 | 5.0 | 0.0 |
| LAS | 3.0 | 3.0 | 3.0 |
| QAS | - | 1.5 | 1.5 |
| DTPMP | 0.4 | 0.2 | 0.4 |
| EDDS | - | 0.4 | 0.2 |
| CMC | 0.4 | 0.4 | 0.4 |
| MA/AA | 4.0 | 2.0 | 2.0 |
| Agglomerates | | | |
| LAS | 5.0 | 5.0 | 5.0 |
| TAS | 2.0 | 2.0 | 1.0 |
| Silicate | 3.0 | 3.0 | 4.0 |
| Zeolite A | 8.0 | 8.0 | 8.0 |
| Carbonate | 8.0 | 8.0 | 4.0 |
| Spray On | | | |
| Perfume | 0.3 | 0.3 | 0.3 |
| C45E7 | 2.0 | 2.0 | 2.0 |
| C25E3 | 2.0 | - | - |
| PARP2 | 0.4 | 0.2 | - |
| Dry additives | | | |
| Citrate | 5.0 | - | 2.0 |
| Bicarbonate | - | 3.0 | - |
| Carbonate | 8.0 | 15.0 | 10.0 |
| TAED | 6.0 | 2.0 | 5.0 |
| PB1 | 14.0 | 7.0 | 10.0 |
| PEO | - | - | 0.2 |

T.01913 "12.01" 8261001

| PARP3 | 1.2(c) | 1.0 (c) | 0.75(c) |
|---------------------|--------|---------|---------|
| Bentonite clay | - | - | 10.0 |
| Protease | 1.0 | 1.0 | 1.0 |
| Lipase | 0.4 | 0.4 | 0.4 |
| Amylase | 0.6 | 0.6 | 0.6 |
| Cellulase | 0.6 | 0.6 | 0.6 |
| Silicone antifoam | 5.0 | 5.0 | 5.0 |
| Sodium sulfate | 0.0 | 3.0 | 0.0 |
| Misc/minors to 100% | 100.0 | 100.0 | 100.0 |
| Density (g/litre) | 850 | 850 | 850 |

| | D | E | F | G | H |
|------------------------------|------|------|------|------|------|
| Blown Powder | | | | | |
| STPP/ Zeolite A | 9.0 | 15.0 | 15.0 | 9.0 | 9.0 |
| Flocculating agent II or III | 0.5 | 0.2 | 0.9 | 1.5 | - |
| LAS | 7.5 | 23.0 | 3.0 | 7.5 | 7.5 |
| QAS | 2.5 | 1.5 | - | - | - |
| DTPMP | 0.4 | 0.2 | 0.4 | 0.4 | 0.4 |
| HEDP or EDDS | - | 0.4 | 0.2 | - | - |
| CMC | 0.1 | 0.4 | 0.4 | 0.1 | 0.1 |
| Sodium carbonate | 5.0 | 20.0 | 20.0 | 10.0 | - |
| Brightener | 0.05 | - | - | 0.05 | 0.05 |
| Clay I or II | - | 10.0 | - | - | - |
| STS | 0.5 | - | - | 0.5 | 0.5 |
| MA/AA | 1.5 | 2.0 | 2.0 | 1.5 | 1.5 |
| Agglomerates | | | | | |
| Suds suppresser (silicon) | 1.0 | 1.0 | - | 2.0 | 0.5 |
| Agglomerate | | | | | |
| Clay | 9.0 | - | - | 4.0 | 10.0 |
| Wax | 0.5 | - | - | 0.5 | 1.5 |
| Glycerol | 0.5 | - | - | 0.5 | 0.5 |
| Agglomerate | | | | | |

| | | | | | |
|----------------------------------|------|------|------|------|------|
| LAS | - | 5.0 | 5.0 | - | - |
| TAS | - | 2.0 | 1.0 | - | - |
| Silicate | - | 3.0 | 4.0 | - | - |
| Zeolite A | - | 8.0 | 8.0 | - | - |
| Carbonate | - | 8.0 | 4.0 | - | - |
| Spray On | | | | | |
| Perfume | - | - | - | - | 0.1 |
| C45E7 or E9 | 2.0 | - | - | 2.0 | 2.0 |
| C25E3 or E4 | 2.0 | - | - | 2.0 | 2.0 |
| Dry additives | | | | | |
| Citrate or citric acid | 2.5 | - | 2.0 | 2.5 | 2.5 |
| Clay I or II | - | 5.0 | 5.0 | - | - |
| Flocculating agent I or II | - | - | - | - | 0.2 |
| Bicarbonate | - | 3.0 | - | - | - |
| Carbonate | 15.0 | - | - | 25.0 | 31.0 |
| TAED | 1.0 | 2.0 | 5.0 | 1.0 | - |
| Sodium perborate or percarbonate | 6.0 | 7.0 | 10.0 | 6.0 | - |
| SRP1, 2, 3 or 4 | 0.2 | 0.1 | 0.2 | 0.5 | 0.3 |
| CMC or nonionic cellulose ether | 1.0 | 1.5 | 0.5 | - | - |
| Protease | 0.3 | 1.0 | 1.0 | 0.3 | 0.3 |
| Lipase | - | 0.4 | 0.4 | - | - |
| Amylase | 0.2 | 0.6 | 0.6 | 0.2 | 0.2 |
| Cellulase | 0.2 | 0.6 | 0.6 | 0.2 | 0.2 |
| Silicone antifoam | - | 5.0 | 5.0 | - | - |
| Perfume (starch) | 0.2 | 0.3 | 1.0 | 0.2 | 0.2 |
| Speckle | 0.5 | 0.5 | 0.1 | - | 1.0 |
| NaSKS-6 (silicate 2R) | 3.5 | - | - | - | 3.5 |
| Photobleach | 0.1 | - | - | 0.1 | 0.1 |
| Soap | 0.5 | 2.5 | - | 0.5 | 0.5 |
| Sodium sulfate | - | 3.0 | - | - | - |
| PARP2 | 0.4 | 0.16 | 0.1 | 0.4 | 0.18 |

| | | | | | | |
|-------------------|----|-------|-------|-------|-------|-------|
| Misc/minors | to | 100.0 | 100.0 | 100.0 | 100.0 | 100.0 |
| 100% | | | | | | |
| Density (g/litre) | | 850 | 850 | 850 | 850 | 850 |

Example 9

The following detergent formulations, according to the present invention were

5 prepared:

| | A | B | C | D |
|----------------|------|------|------|------|
| LAS | 18.0 | 14.0 | 24.0 | 20.0 |
| QAS | 0.7 | 1.0 | - | 0.7 |
| TFAA | - | 1.0 | - | - |
| C23E56.5 | - | - | 1.0 | - |
| C45E7 | - | 1.0 | - | - |
| C45E3S | 1.0 | 2.5 | 1.0 | - |
| STPP | 32.0 | 18.0 | 30.0 | 22.0 |
| Silicate | 9.0 | 5.0 | 9.0 | 8.0 |
| Carbonate | 9.0 | 7.5 | - | 5.0 |
| Bicarbonate | - | 7.5 | - | - |
| PB1 | 3.0 | 1.0 | - | - |
| PB4 | - | 1.0 | - | - |
| NOBS | 2.0 | 1.0 | - | - |
| DTPMP | - | 1.0 | - | - |
| DTPA | 0.5 | - | 0.2 | 0.3 |
| SRP 1 | 0.3 | 0.2 | - | 0.1 |
| MA/AA | 1.0 | 1.5 | 2.0 | 0.5 |
| CMC | 0.8 | 0.4 | 0.4 | 0.2 |
| PEI | - | - | 0.4 | - |
| Sodium sulfate | 20.0 | 10.0 | 20.0 | 30.0 |
| Mg sulfate | 0.2 | - | 0.4 | 0.9 |
| Protease | 0.8 | 1.0 | 0.5 | 0.5 |
| Amylase | 0.5 | 0.4 | - | 0.25 |
| Lipase | 0.2 | - | 0.1 | - |

| | | | | |
|-----------------------------|--------|--------|--------|--------|
| Cellulase | 0.15 | - | - | 0.05 |
| Photoactivated bleach (ppm) | 30ppm | 20ppm | - | 10ppm |
| PARP3 | 0.2(c) | 1.4(s) | 0.8(c) | 0.8(c) |
| Perfume spray on | 0.3 | 0.3 | 0.1 | 0.2 |
| Brightener 1/2 | 0.05 | 0.2 | 0.08 | 0.1 |
| Misc/minors to 100% | | | | |

Example 10

The following is a composition in the form of a tablet, bar, extrudate or granule in accord with the invention

| | A | B | C | D | E | F | G |
|---|------|------|------|------|------|------|------|
| Sodium C ₁₁ -C ₁₃ alkylbenzenesulfonate | 12.0 | 16.0 | 23.0 | 19.0 | 18.0 | 20.0 | 16.0 |
| Sodium C ₁₄ -C ₁₅ alcohol sulfate | | 4.5 | - | | - | - | 4.0 |
| C ₁₄ -C ₁₅ alcohol ethoxylate (3) sulfate | - | - | 2.0 | - | 1.0 | 1.0 | 1.0 |
| Sodium C ₁₄ -C ₁₅ alcohol ethoxylate | 2.0 | 2.0 | - | 1.3 | - | - | 5.0 |
| C ₉ -C ₁₄ alkyl dimethyl hydroxy ethyl quaternary ammonium salt | | | - | - | 1.0 | 0.5 | 2.0 |
| Tallow fatty acid | | | - | - | - | - | 1.0 |
| Sodium tripolyphosphate / Zeolite | 23.0 | 25.0 | 14.0 | 22.0 | 20.0 | 10.0 | 20.0 |
| Sodium carbonate | 25.0 | 22.0 | 35.0 | 20.0 | 28.0 | 41.0 | 30.0 |
| Sodium Polyacrylate (45%) | 0.5 | 0.5 | 0.5 | 0.5 | - | - | - |
| Sodium polyacrylate/maleate polymer | - | - | 1.0 | 1.0 | 1.0 | 2.0 | 0.5 |

| | | | | | | | |
|--|-----|-----|------|------|------|-----|-----|
| Sodium silicate (1:6 ratio NaO/SiO ₂)(46%) | 3.0 | 6.0 | 9.0 | 8.0 | 9.0 | 6.0 | 8.0 |
| Sodium sulfate | - | - | - | - | - | 2.0 | 3.0 |
| Sodium perborate/ percarbonate | 5.0 | 5.0 | 10.0 | - | 3.0 | 1.0 | - |
| Poly(ethyleneglycol), MW ~4000 (50%) | 1.5 | 1.5 | 1.0 | 1.0 | - | - | 0.5 |
| Sodium carboxy methyl cellulose | 1.0 | 1.0 | 1.0 | - | 0.5 | 0.5 | 0.5 |
| NOBS/ DOBS | - | 1.0 | - | - | 1.0 | 0.7 | - |
| TAED | 1.5 | 1.0 | 2.5 | - | 3.0 | 0.7 | - |
| SRP 1 | 1.5 | 1.5 | 1.0 | 1.0 | - | 1.0 | - |
| Clay I or II | 5.0 | 6.0 | 12.0 | 7.0 | 10.0 | 4.0 | 3.0 |
| Flocculating agent I or III | 0.2 | 0.2 | 3.0 | 2.0 | 0.1 | 1.0 | 0.5 |
| Humectant | 0.5 | 1.0 | 0.5 | 1.0 | 0.5 | 0.5 | - |
| Wax | 0.5 | 0.5 | 1.0 | - | - | 0.5 | 0.5 |
| Moisture | 7.5 | 7.5 | 6.0 | 7.0 | 5.0 | 3.0 | 5.0 |
| Magnesium sulphate | - | - | - | - | - | 0.5 | 1.5 |
| Chelant | - | - | - | - | 0.8 | 0.6 | 1.0 |
| Enzymes, including amylase, cellulase, protease and lipase | - | - | - | - | 2.0 | 1.5 | 2.0 |
| Speckle | 2.5 | 4.1 | 4.2 | 4.4 | 5.6 | 5.0 | 5.2 |
| minors, e.g. PVP, PVPVI/PVNO, brightener, photo-bleach, | 2.0 | 1.0 | 1.0 | 1.0 | 2.5 | 1.5 | 1.0 |
| PARP2 | 0.4 | 0.8 | 0.6 | 0.2 | 0.1 | 0.8 | 0.1 |
| Perfume | 0.1 | - | - | 0.05 | 0.2 | - | 0.3 |

| | H | I | J | K |
|--|------|------|------|------|
| Sodium C ₁₁ -C ₁₃ alkylbenzenesulfonate | 23.0 | 13.0 | 20.0 | 18.0 |
| Sodium C ₁₄ -C ₁₅ alcohol sulfate | - | 4.0 | - | - |
| Clay I or II | 5.0 | 10.0 | 14.0 | 6.0 |
| Flocculating agent I or II | 0.2 | 0.3 | 0.1 | 0.9 |

| | | | | |
|--|------|------|------|------|
| Wax | 0.5 | 0.5 | 1.0 | - |
| Humectant (glycerol/ silica) | 0.5 | 2.0 | 1.5 | - |
| C ₁₄ -C ₁₅ alcohol ethoxylate sulfate | - | | - | 2.0 |
| Sodium C ₁₄ -C ₁₅ alcohol ethoxylate (| 2.5 | 3.5 | - | - |
| C ₉ -C ₁₄ alkyl dimethyl hydroxy ethyl quaternary ammonium salt | | - | - | 0.5 |
| Tallow fatty acid | 0.5 | - | - | - |
| Tallow alcohol ethoxylate (50) | - | - | | 1.3 |
| Sodium tripolyphosphate | - | 41.0 | - | 20.0 |
| Zeolite A, hydrate (0.1-10 micron size) | 26.3 | - | 21.3 | - |
| Sodium carbonate | 24.0 | 22.0 | 35.0 | 27.0 |
| Sodium Polyacrylate (45%) | 2.4 | - | 2.7 | - |
| Sodium polyacrylate/maleate polymer | - | - | 1.0 | 2.5 |
| Sodium silicate (1.6 or 2 or 2.2 ratio NaO/SiO ₂)(46%) | 4.0 | 7.0 | 2.0 | 6.0 |
| Sodium sulfate | - | 6.0 | 2.0 | - |
| Sodium perborate/ percarbonate | 8.0 | 4.0 | - | 12.0 |
| Poly(ethyleneglycol), MW ~4000 (50%) | 1.7 | 0.4 | 1.0 | - |
| Sodium carboxy methyl cellulose | 1.0 | - | - | 0.3 |
| Citric acid | - | - | 3.0 | - |
| NOBS/ DOBS | 1.2 | - | - | 1.0 |
| TAED | 0.6 | 1.5 | - | 3.0 |
| Perfume | 0.5 | 0.1 | 0.3 | - |
| SRP 1 | - | 1.5 | 1.0 | 1.0 |
| Moisture | 7.5 | 3.1 | 6.1 | 7.3 |
| Magnesium sulphate | - | - | - | 1.0 |
| Chelant | - | - | - | 0.5 |
| speckle | 1.0 | 0.5 | 0.2 | 2.7 |
| Enzymes, including amylase, cellulase, protease and lipase | - | 1.0 | - | 1.5 |
| minors, e.g. brightener, photo-bleach | 1.0 | 1.0 | 1.0 | 1.0 |
| PARP2 | 0.2 | 0.4 | 0.16 | 0.8 |

Example 11

The following liquid detergent formulations were prepared in accord with the invention (levels are given as parts per weight).

| | A | B | C | D | E |
|-------------------------|--------|--------|--------|------|--------|
| LAS | 11.5 | 8.8 | - | 3.9 | - |
| C25E2.5S | - | 3.0 | 18.0 | - | 16.0 |
| C45E2.25S | 11.5 | 3.0 | - | 15.7 | - |
| C23E9 | - | 2.7 | 1.8 | 2.0 | 1.0 |
| C23E7 | 3.2 | - | - | - | - |
| CFAA | - | - | 5.2 | - | 3.1 |
| TPKFA | 1.6 | - | 2.0 | 0.5 | 2.0 |
| Citric acid (50%) | 6.5 | 1.2 | 2.5 | 4.4 | 2.5 |
| Calcium formate | 0.1 | 0.06 | 0.1 | - | - |
| Sodium formate | 0.5 | 0.06 | 0.1 | 0.05 | 0.05 |
| Sodium cumene sulfonate | 4.0 | 1.0 | 3.0 | 1.18 | - |
| Borate | 0.6 | - | 3.0 | 2.0 | 2.9 |
| Sodium hydroxide | 5.8 | 2.0 | 3.5 | 3.7 | 2.7 |
| Ethanol | 1.75 | 1.0 | 3.6 | 4.2 | 2.9 |
| 1, 2 propanediol | 3.3 | 2.0 | 8.0 | 7.9 | 5.3 |
| Monoethanolamine | 3.0 | 1.5 | 1.3 | 2.5 | 0.8 |
| TEPAE | 1.6 | - | 1.3 | 1.2 | 1.2 |
| Protease | 1.0 | 0.3 | 1.0 | 0.5 | 0.7 |
| Lipase | - | - | 0.1 | - | - |
| Cellulase | - | - | 0.1 | 0.2 | 0.05 |
| Amylase | - | - | - | 0.1 | - |
| SRP1 | 0.2 | - | 0.1 | - | - |
| DTPA | - | - | 0.3 | - | - |
| PVNO | - | - | 0.3 | - | 0.2 |
| PARP1 | 0.8(c) | 1.0(c) | - | - | - |
| PARP3 | - | - | 0.3(c) | 0.4 | 0.4(s) |
| Brightener 1 | 0.2 | 0.07 | 0.1 | - | - |
| Silicone antifoam | 0.04 | 0.02 | 0.1 | 0.1 | 0.1 |
| Water/minors up to 100% | | | | | |

The following liquid detergent formulations were prepared in accord with the invention (levels are given in parts per weight):

| | A | B | C | D | E | F | G | H |
|--------------------------------------|------|------|------|------|------|------|------|------|
| LAS | 10.0 | 13.0 | 9.0 | - | 25.0 | - | - | - |
| C25AS | 4.0 | 1.0 | 2.0 | 10.0 | - | 13.0 | 18.0 | 15.0 |
| C25E3S | 1.0 | - | - | 3.0 | - | 2.0 | 2.0 | 4.0 |
| C25E7 | 6.0 | 8.0 | 13.0 | 2.5 | - | - | 4.0 | 4.0 |
| TFAA | - | - | - | 4.5 | - | 6.0 | 8.0 | 8.0 |
| APA | - | 1.4 | - | - | 3.0 | 1.0 | 2.0 | - |
| TPKFA | 2.0 | - | 13.0 | 7.0 | - | 15.0 | 11.0 | 11.0 |
| Citric acid | 2.0 | 3.0 | 1.0 | 1.5 | 1.0 | 1.0 | 1.0 | 1.0 |
| Dodecenyl/tetradecenyl succinic acid | 12.0 | 10.0 | - | - | 15.0 | - | - | - |
| Rape seed fatty acid | 4.0 | 2.0 | 1.0 | - | 1.0 | - | 3.5 | - |
| Ethanol | 4.0 | 4.0 | 7.0 | 2.0 | 7.0 | 2.0 | 3.0 | 2.0 |
| 1,2 Propanediol | 4.0 | 4.0 | 2.0 | 7.0 | 6.0 | 8.0 | 10.0 | 13.0 |
| Monoethanolamine | - | - | - | 5.0 | - | - | 9.0 | 9.0 |
| Triethanolamine | - | - | 8.0 | - | - | - | - | - |
| TEPAE | 0.5 | - | 0.5 | 0.2 | - | - | 0.4 | 0.3 |
| DTPMP | 1.0 | 1.0 | 0.5 | 1.0 | 2.0 | 1.2 | 1.0 | - |
| Protease | 0.5 | 0.5 | 0.4 | 0.25 | - | 0.5 | 0.3 | 0.6 |
| Alcalase | - | - | - | - | 1.5 | - | - | - |
| Lipase | - | 0.10 | - | 0.01 | - | - | 0.15 | 0.15 |
| Amylase | 0.25 | 0.25 | 0.6 | 0.5 | 0.25 | 0.9 | 0.6 | 0.6 |
| Cellulase | - | - | - | 0.05 | - | - | 0.15 | 0.15 |
| Endolase | - | - | - | 0.10 | - | - | 0.07 | - |
| SRP2 | 0.3 | - | 0.3 | 0.1 | - | - | 0.2 | 0.1 |
| Boric acid | 0.1 | 0.2 | 1.0 | 2.0 | 1.0 | 1.5 | 2.5 | 2.5 |
| Calcium chloride | - | 0.02 | - | 0.01 | - | - | - | - |
| Bentonite clay | - | - | - | - | 4.0 | 4.0 | - | - |
| Brightener 1 | - | 0.4 | - | - | 0.1 | 0.2 | 0.3 | - |
| Sud supressor | 0.1 | 0.3 | - | 0.1 | 0.4 | - | - | - |
| Opacifier | 0.5 | 0.4 | - | 0.3 | 0.8 | 0.7 | - | - |

| | | | | | | | | |
|-------------------------|------------|------------|------------|------------|------------|------------|------------|-------------|
| PARP1 | 0.8 (s) | - | 0.5 (c) | - | 0.3 (c) | - | 0.4 (c) | 0.2 (c) |
| PARP3 | - | 0.2 (c) | - | 0.1 (c) | - | 0.8 (c) | 0.1 (c) | 0.07 (c) |
| Water/minors up to 100% | | | | | | | | |
| NaOH up to pH | 8.0 | 8.0 | 7.6 | 7.7 | 8.0 | 7.5 | 8.0 | 8.2 |

Example 13

The following liquid detergent compositions were prepared in accord with the invention (levels are given in parts per weight).

| | A | B |
|-------------------------|--------|--------|
| LAS | 27.6 | 18.9 |
| C45AS | 13.8 | 5.9 |
| C13E8 | 3.0 | 3.1 |
| Oleic acid | 3.4 | 2.5 |
| Citric acid | 5.4 | 5.4 |
| Sodium hydroxide | 0.4 | 3.6 |
| Calcium formate | 0.2 | 0.1 |
| Sodium formate | - | 0.5 |
| Ethanol | 7.0 | - |
| Monoethanolamine | 16.5 | 8.0 |
| 1,2 propanediol | 5.9 | 5.5 |
| Xylene sulfonic acid | - | 2.4 |
| TEPAE | 1.5 | 0.8 |
| Protease | 1.5 | 0.6 |
| PEG | - | 0.7 |
| Brightener 2 | 0.4 | 0.1 |
| Perfume spray on | 0.5 | 0.3 |
| PARP3 | 0.2(c) | - |
| PARP1 | 0.1(c) | 0.2(c) |
| Water/minors up to 100% | | |

Example 14

The following laundry bar detergent compositions were prepared in accord with the invention (levels are given in parts per weight).

5

| | A | B | C | D | E | F |
|-------------------|------|------|------|--------|------|------|
| LAS | - | - | 19.0 | 6.75 | 8.8 | - |
| C28AS | 30.0 | 13.5 | - | 15.75 | 11.2 | 22.5 |
| Sodium laurate | 2.5 | 9.0 | - | - | - | - |
| Zeolite A | 2.0 | 1.25 | - | 1.25 | 1.25 | 1.25 |
| Carbonate | 10.0 | - | 11.0 | 7.0 | 13.0 | 9.0 |
| Calcium carbonate | 27.5 | 39.0 | 35.0 | 40.0 | - | 40.0 |
| Sulfate | 5.0 | 5.0 | 3.0 | - | - | 5.0 |
| TSP | 5.0 | - | - | 5.0 | 2.5 | - |
| STPP | 5.0 | 15.0 | 10.0 | 7.0 | 8.0 | 10.0 |
| Bentonite clay | - | 10.0 | - | - | - | - |
| DTPMP | - | 0.7 | 0.6 | 0.7 | 0.7 | 0.7 |
| CMC | - | 1.0 | 1.0 | - | - | 1.0 |
| Talc | - | - | 10.0 | - | - | - |
| Silicate | - | - | 4.0 | - | - | - |
| PVNO | 0.02 | 0.03 | - | 0.02 | - | - |
| MA/AA | 0.4 | 1.0 | - | 0.4 | 0.5 | 0.4 |
| SRP1 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 |
| Protease | - | 0.12 | - | - | - | 0.1 |
| Lipase | - | 0.1 | - | - | - | - |
| Amylase | - | - | 0.8 | - | 0.1 | - |
| Cellulase | - | 0.15 | - | 0.1 | - | - |
| PEO | - | 0.2 | - | - | - | 0.3 |
| Perfume | 0.2 | 0.5 | 0.3 | - | - | 0.05 |
| Mg sulfate | - | - | 3.0 | - | - | - |
| PARP4 | 0.8 | - | - | 0.8(c) | - | - |
| PARP2 | - | 0.2 | - | - | 0.8 | - |
| PARP3 | - | - | 1(c) | - | - | 1(c) |
| Brightener | 0.15 | 0.10 | 0.15 | - | - | 0.1 |

| | | | | | | |
|--------------------------------|---|------|------|---|---|------|
| Photoactivated bleach (ppm) | - | 15.0 | 15.0 | - | - | 15.0 |
|--------------------------------|---|------|------|---|---|------|

Example 15

The following detergent additive compositions were prepared according to the present invention :

| | A | B | C |
|---------------------------------|----------|------------|----------|
| LAS | - | 5.0 | 5.0 |
| STPP | 30.0 | - | 20.0 |
| Zeolite A | - | 35.0 | 20.0 |
| PB1 | 20.0 | 15.0 | - |
| TAED | 10.0 | 8.0 | - |
| PARP1 | 1.0(c) | - | 0.5(c) |
| PARP3 | - | 0.3(c) | 0.1(c) |
| Protease | - | 0.3 | 0.3 |
| Amylase | - | 0.06 | 0.06 |
| Minors, water and miscellaneous | | Up to 100% | |

Example 16

The following compact high density (0.96Kg/l) dishwashing detergent compositions were prepared according to the present invention :

| | A | B | C | D | E | F | G | H |
|--------------|----------|----------|----------|----------|----------|----------|----------|----------|
| STPP | - | - | 54.3 | 51.4 | 51.4 | - | - | 50.9 |
| Citrate | 35.0 | 17.0 | - | - | - | 46.1 | 40.2 | - |
| Carbonate | - | 15.0 | 12.0 | 14.0 | 4.0 | - | 7.0 | 31.1 |
| Bicarbonate | - | - | - | - | - | 25.4 | - | - |
| Silicate | 32.0 | 14.8 | 14.8 | 10.0 | 10.0 | 1.0 | 25.0 | 3.1 |
| Metasilicate | - | 2.5 | - | 9.0 | 9.0 | - | - | - |
| PB1 | 1.9 | 9.7 | 7.8 | 7.8 | 7.8 | - | - | - |
| PB4 | 8.6 | - | - | - | - | - | - | - |
| Percarbonate | - | - | - | - | - | 6.7 | 11.8 | 4.8 |

| | | | | | | | | |
|-------------------------|------------|--------|--------|--------|--------|--------|--------|--------|
| Nonionic | 1.5 | 2.0 | 1.5 | 1.7 | 1.5 | 2.6 | 1.9 | 5.3 |
| TAED | 5.2 | 2.4 | - | - | - | 2.2 | - | 1.4 |
| HEDP | - | 1.0 | - | - | - | - | - | - |
| DTPMP | - | 0.6 | - | - | - | - | - | - |
| MnTACN | - | - | - | - | - | - | 0.008 | - |
| PAAC | - | - | 0.008 | 0.01 | 0.007 | - | - | - |
| BzP | - | - | - | - | 1.4 | - | - | - |
| Paraffin | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | - | - |
| PARP3 | 0.2 | 0.4(c) | 0.2(c) | 0.1(c) | - | - | - | 0.5(c) |
| PARP1 | - | - | - | - | 1.0(c) | 1.3(c) | 1.2(c) | - |
| Protease | 0.072 | 0.072 | 0.029 | 0.053 | 0.046 | 0.026 | 0.059 | 0.06 |
| Amylase | 0.012 | 0.012 | 0.006 | 0.012 | 0.013 | 0.009 | 0.017 | 0.03 |
| Lipase | - | 0.001 | - | 0.005 | - | - | - | - |
| BTA | 0.3 | 0.3 | 0.3 | 0.3 | 0.3 | - | 0.3 | 0.3 |
| MA/AA | - | - | - | - | - | - | 4.2 | - |
| 480N | 3.3 | 6.0 | - | - | - | - | - | 0.9 |
| Perfume | 0.2 | 0.2 | 0.2 | 0.2 | 0.05 | - | - | 0.1 |
| Sulphate | 7.0 | 20.0 | 5.0 | 2.2 | 0.8 | 12.0 | 4.6 | - |
| pH | 10.8 | 11.0 | 10.8 | 11.3 | 11.3 | 9.6 | 10.8 | 10.9 |
| Miscellaneous and water | Up to 100% | | | | | | | |

Example 17

- 5 The following granular dishwashing detergent compositions of bulk density 1.02Kg/L were prepared according to the present invention :

| | A | B | C | D | E | F | G | H |
|--------------|------|------|------|------|------|------|------|------|
| STPP | 30.0 | 30.0 | 33.0 | 34.2 | 29.6 | 31.1 | 26.6 | 17.6 |
| Carbonate | 29.5 | 30.0 | 29.0 | 24.0 | 15.0 | 36.0 | 2.1 | 38.0 |
| Silicate | 7.4 | 7.4 | 7.5 | 7.2 | 13.3 | 3.4 | 43.7 | 12.4 |
| Metasilicate | - | - | 4.5 | 5.1 | - | - | - | - |
| Percarbonate | - | - | - | - | - | 4.0 | - | - |
| PB1 | 4.4 | 4.2 | 4.5 | 4.5 | - | - | - | - |
| NADCC | - | - | - | - | 2.0 | - | 1.6 | 1.0 |
| Nonionic | 1.2 | 1.0 | 0.7 | 0.8 | 1.9 | 0.7 | 0.6 | 0.3 |

| | | | | | | | | |
|-------------------------|--------|--------|--------|--------|------------|--------|--------|--------|
| TAED | 1.0 | - | - | - | - | 0.8 | - | - |
| PAAC | - | 0.004 | 0.004 | 0.004 | - | - | - | - |
| BzP | - | - | - | 1.4 | - | - | - | - |
| Paraffin | 0.25 | 0.25 | 0.25 | 0.25 | - | - | - | - |
| PARP3 | 0.1(c) | 0.5(c) | 0.4(c) | 0.8(c) | - | - | 1.0(c) | 0.5(c) |
| PARP1 | - | - | - | - | 1.0 | 1.5(c) | 1.2(c) | 0.8 |
| | | | | | (c) | | | (s) |
| Protease | 0.036 | 0.015 | 0.03 | 0.028 | - | 0.03 | - | - |
| Amylase | 0.003 | 0.003 | 0.01 | 0.006 | - | 0.01 | - | - |
| Lipase | 0.005 | - | 0.001 | - | - | - | - | - |
| BTA | 0.15 | 0.15 | 0.15 | 0.15 | - | - | - | - |
| Perfume | 0.2 | 0.2 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | - |
| Sulphate | 23.4 | 25.0 | 22.0 | 18.5 | 30.1 | 19.3 | 23.1 | 23.6 |
| pH | 10.8 | 10.8 | 11.3 | 11.3 | 10.7 | 11.5 | 12.7 | 10.9 |
| Miscellaneous and water | | | | | Up to 100% | | | |

Example 18

- 5 The following tablet detergent compositions were prepared according to the present invention by compression of a granular dishwashing detergent composition at a pressure of 13KN/cm² using a standard 12 head rotary press:

| | A | B | C | D | E | F |
|-----------|--------|-------|--------|--------|-------|-------|
| STPP | - | 48.8 | 49.2 | 38.0 | - | 46.8 |
| Citrate | 26.4 | - | - | - | 31.1 | - |
| Carbonate | - | 4.0 | 12.0 | 14.4 | 10.0 | 20.0 |
| Silicate | 26.4 | 14.8 | 15.0 | 12.6 | 17.7 | 2.4 |
| PARP1 | 1.3(c) | - | - | - | 1.5 | - |
| PARP2 | - | 0.2 | - | - | - | 0.4 |
| PARP3 | - | - | 0.2(c) | 0.1(s) | - | - |
| Protease | 0.058 | 0.072 | 0.041 | 0.033 | 0.052 | 0.013 |
| Amylase | 0.01 | 0.03 | 0.012 | 0.007 | 0.016 | 0.002 |
| Lipase | 0.005 | - | - | - | - | - |
| PB1 | 1.6 | 7.7 | 12.2 | 10.6 | 15.7 | - |
| PB4 | 6.9 | - | - | - | - | 14.4 |
| Nonionic | 1.5 | 2.0 | 1.5 | 1.65 | 0.8 | 6.3 |

70

| | | | | | | |
|-------------------------|------------|------|------|-------|-------|------|
| PAAC | - | - | 0.02 | 0.009 | - | - |
| MnTACN | - | - | - | - | 0.007 | - |
| TAED | 4.3 | 2.5 | - | - | 1.3 | 1.8 |
| HEDP | 0.7 | - | - | 0.7 | - | 0.4 |
| DTPMP | 0.65 | - | - | - | - | - |
| Paraffin | 0.4 | 0.5 | 0.5 | 0.55 | - | - |
| BTA | 0.2 | 0.3 | 0.3 | 0.3 | - | - |
| PA30 | 3.2 | - | - | - | - | - |
| MA/AA | - | - | - | - | 4.5 | 0.55 |
| Perfume | - | - | 0.05 | 0.05 | 0.2 | 0.2 |
| Sulphate | 24.0 | 13.0 | 2.3 | - | 10.7 | 3.4 |
| Weight of tablet | 25g | 25g | 20g | 30g | 18g | 20g |
| pH | 10.6 | 10.6 | 10.7 | 10.7 | 10.9 | 11.2 |
| Miscellaneous and water | Up to 100% | | | | | |

Example 19

The following liquid dishwashing detergent compositions of density 1.40Kg/L were prepared according to the present invention :

5

| | A | B | C | D |
|-----------------------------------|------------|----------|----------|----------|
| STPP | 17.5 | 17.5 | 17.2 | 16.0 |
| Carbonate | 2.0 | - | 2.4 | - |
| Silicate | 5.3 | 6.1 | 14.6 | 15.7 |
| NaOCl | 1.15 | 1.15 | 1.15 | 1.25 |
| Polygen/carbopol | 1.1 | 1.0 | 1.1 | 1.25 |
| Nonionic | - | - | 0.1 | - |
| NaBz | 0.75 | 0.75 | - | - |
| PARP3 | 0.4(c) | 0.2(c) | 0.1(c) | 0.5(c) |
| NaOH | - | 1.9 | - | 3.5 |
| KOH | 2.8 | 3.5 | 3.0 | - |
| pH | 11.0 | 11.7 | 10.9 | 11.0 |
| Sulphate, miscellaneous and water | up to 100% | | | |

10 **Example 20**

The following liquid rinse aid compositions were prepared according to the present invention :

| | A | B | C |
|-------------------------|--------|------------|--------|
| Nonionic | 12.0 | - | 14.5 |
| Nonionic blend | - | 64.0 | - |
| Citric | 3.2 | - | 6.5 |
| HEDP | 0.5 | - | - |
| PEG | - | 5.0 | - |
| SCS | 4.8 | - | 7.0 |
| Ethanol | 6.0 | 8.0 | - |
| PARP1 | 1.6(c) | - | 1.0(c) |
| PARP3 | - | 1.0(c) | 0.1(c) |
| pH of the liquid | 2.0 | 7.5 | / |
| Miscellaneous and water | | Up to 100% | |

Example 21

The following liquid dishwashing compositions were prepared according to the present invention :

| | A | B | C | D | E |
|---------------------------------------|------|------|------|------|------|
| C17ES | 28.5 | 27.4 | 19.2 | 34.1 | 34.1 |
| Amine oxide | 2.6 | 5.0 | 2.0 | 3.0 | 3.0 |
| C12 glucose amide | - | - | 6.0 | - | - |
| Betaine | 0.9 | - | - | 2.0 | 2.0 |
| Xylene sulfonate | 2.0 | 4.0 | - | 2.0 | - |
| Neodol C11E9 | - | - | 5.0 | - | - |
| Polyhydroxy fatty acid amide | - | - | - | 6.5 | 6.5 |
| Sodium diethylene penta acetate (40%) | - | - | 0.03 | - | - |
| TAED | - | - | - | 0.06 | 0.06 |
| Sucrose | - | - | - | 1.5 | 1.5 |
| Ethanol | 4.0 | 5.5 | 5.5 | 9.1 | 9.1 |
| Alkyl diphenyl oxide disulfonate | - | - | - | - | 2.3 |
| Ca formate | - | - | - | 0.5 | 1.1 |

72

| | | | | | |
|-------------------|------------|--------|--------|--------|---------|
| Ammonium citrate | 0.06 | 0.1 | - | - | - |
| Na chloride | - | 1.0 | - | - | - |
| Mg chloride | 3.3 | - | 0.7 | - | - |
| Ca chloride | - | - | 0.4 | - | - |
| Na sulfate | - | - | 0.06 | - | - |
| Mg sulfate | 0.08 | - | - | - | - |
| Mg hydroxide | - | - | - | 2.2 | 2.2 |
| Na hydroxide | - | - | - | 1.1 | 1.1 |
| Hydrogen peroxide | 200ppm | 0.16 | 0.006 | - | - |
| PARP3 | 0.4(c) | - | 0.2(c) | - | 0.25(c) |
| PARP1 | - | 1.6(c) | - | 1.4(s) | 1.3(c) |
| Protease | 0.017 | 0.005 | .0035 | 0.003 | 0.002 |
| Perfume | 0.18 | 0.09 | 0.09 | 0.2 | 0.2 |
| Water and minors | Up to 100% | | | | |

Example 22

- 5 The following liquid hard surface cleaning compositions were prepared according to the present invention :

| | A | B | C | D | E |
|-------------------------|--------|--------|--------|--------|---------|
| PARP1 | 1.8(c) | - | 1.6(s) | - | 1.4(c) |
| PARP3 | - | 0.2(c) | - | 0.1(c) | 0.05(c) |
| Amylase | 0.01 | 0.002 | 0.005 | - | - |
| Protease | 0.05 | 0.01 | 0.02 | - | - |
| Hydrogen peroxide | - | - | - | 6.0 | 6.8 |
| Acetyl triethyl citrate | - | - | - | 2.5 | - |
| DTPA | - | - | - | 0.2 | - |
| Butyl hydroxy toluene | - | - | - | 0.05 | - |
| EDTA* | 0.05 | 0.05 | 0.05 | - | - |
| Citric / Citrate | 2.9 | 2.9 | 2.9 | 1.0 | - |
| LAS | 0.5 | 0.5 | 0.5 | - | - |
| C12 AS | 0.5 | 0.5 | 0.5 | - | - |
| C10AS | - | - | - | - | 1.7 |
| C12(E)S | 0.5 | 0.5 | 0.5 | - | - |
| C12,13 E6.5 nonionic | 7.0 | 7.0 | 7.0 | - | - |

73

| | | | | | |
|---------------------------------------|------|------|------|------------|-----|
| Neodol 23-6.5 | - | - | - | 12.0 | - |
| Dobanol 23-3 | - | - | - | - | 1.5 |
| Dobanol 91-10 | - | - | - | - | 1.6 |
| C25AE1.8S | - | - | - | 6.0 | |
| Na paraffin sulphonate | - | - | - | 6.0 | |
| Perfume | 1.0 | 1.0 | 1.0 | 0.5 | 0.2 |
| Propanediol | - | - | - | 1.5 | |
| Ethoxylated tetraethylene pentaime | - | - | - | 1.0 | - |
| 2, Butyl octanol | - | - | - | - | 0.5 |
| Hexyl carbitol** | 1.0 | 1.0 | 1.0 | - | - |
| SCS | 1.3 | 1.3 | 1.3 | - | - |
| pH adjusted to | 7-12 | 7-12 | 7-12 | 4 | - |
| Miscellaneous and water | | | | Up to 100% | |
| *Na4 ethylenediamine diacetic acid | | | | | |
| **Diethylene glycol monohexyl ether | | | | | |

5

Example 23

The following spray composition for cleaning of hard surfaces and removing household mildew was prepared according to the present invention :

| | |
|--------------------|------------|
| PARP3 | 0.4(c) |
| Amylase | 0.01 |
| Protease | 0.01 |
| Na octyl sulfate | 2.0 |
| Na dodecyl sulfate | 4.0 |
| Na hydroxide | 0.8 |
| Silicate | 0.04 |
| Butyl carbitol* | 4.0 |
| Perfume | 0.35 |
| Water/minors | up to 100% |

10 *Diethylene glycol monobutyl ether

Example 24

The following lavatory cleansing block compositions were prepared according to the present invention.

| | A | B | C |
|---|--------|---------|---------|
| C16-18 fatty alcohol/50EO | 70.0 | - | - |
| LAS | - | - | 80.0 |
| Nonionic | - | 1.0 | - |
| Oleoamide surfactant | - | 25.0 | - |
| Partially esterified copolymer of vinylmethyl ether and maleic anhydride, viscosity 0.1-0.5 | 5.0 | - | - |
| Polyethylene glycol MW 8000 | - | 38.0 | - |
| Water-soluble K-polyacrylate MW 4000-8000 | - | 12.0 | - |
| Water-soluble Na-copolymer of acrylamide (70%) and acrylic acid (30%) low MW | - | 19.0 | - |
| Na triphosphate | 10.0 | - | - |
| Carbonate | - | - | - |
| PARP1 | 0.8(c) | - | 1.6(c) |
| PARP3 | - | 0.2(c) | 0.05(c) |
| Dye | 2.5 | 1.0 | 1.0 |
| Perfume | 3.0 | - | 7.0 |
| KOH / HCL solution | | pH 6-11 | |

5

Example 25

The following toilet bowl cleaning composition was prepared according to the present invention.

| | A | B |
|---------------------------|--------|--------|
| C14-15 linear alcohol 7EO | 2.0 | 10.0 |
| Citric acid | 10.0 | 5.0 |
| PARP1 | 1.2(c) | - |
| PARP3 | - | 0.4(c) |
| DTPMP | - | 1.0 |
| Dye | 2.0 | 1.0 |
| Perfume | 3.0 | 3.0 |

10

NaOH

pH 6-11

Water and minors

Up to 100%

Example 26

- 5 The following fabric softening compositions are in accordance with the present invention

| Component | A | B | C | D | E | F |
|-------------------------------------|---------|--------|--------|--------|--------|--------|
| DTDMAC | - | - | - | - | 4.5 | 15.0 |
| DEQA | 2.6 | 2.9 | 18.0 | 19.0 | - | - |
| Fatty acid | 0.3 | - | 1.0 | - | - | - |
| HCl | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 | 0.02 |
| PEG | - | - | 0.6 | 0.6 | - | 0.6 |
| Perfume | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 |
| Silicone antifoam | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 | 0.01 |
| PARP 3 | 0.45(c) | 0.1(c) | 0.3(s) | 0.3(c) | 0.1(s) | 0.5(c) |
| Electrolyte (ppm) | - | - | 600 | 1200 | - | 1200 |
| Dye (ppm) | 10 | 10 | 50 | 50 | 10 | 50 |
| Water and minors to balance to 100% | | | | | | |

10 **Example 27**

The following dryer added fabric conditioner compositions were prepared according to the present invention :

| | A | B | C | D |
|--------------|------|------|------|------|
| DEQA(2) | - | - | - | 50.0 |
| DTMAMS | - | - | 26.0 | - |
| SDASA | 70.0 | 70.0 | 42.0 | 35.0 |
| Neodol 45-13 | 13.0 | 13.0 | - | - |
| Ethanol | 1.0 | 1.0 | - | - |
| PARP 3(c) | 1.5 | - | 1.5 | 3.0 |
| PARP 3(s) | - | 1.5 | - | - |
| Perfume | 0.75 | 0.75 | 1.0 | 1.5 |

76

| | | | | |
|---------------------------|------|------|------|------|
| Glycoperse S-20 | - | - | - | 10.0 |
| Glycerol monostearate | - | - | 26.0 | - |
| Digeranyl Succinate | 0.38 | 0.38 | - | - |
| Clay | - | - | 3.0 | - |
| Dye | 0.01 | 0.01 | - | - |
| Minors to balance to 100% | | | | |

Example 28

The following are non-limiting examples of pre-soak fabric conditioning and/or fabric enhancement compositions according to the present invention which can be suitably used in the laundry rinse cycle.

5

| Ingredients | A | B | C | D | E | F |
|---|---------|---------|---------|---------|---------|---------|
| Polymer | 3.5 | 3.5 | 3.5 | 3.5 | 3.5 | 3.5 |
| Dye fixative | 2.3 | 2.3 | 2.4 | 2.4 | 2.5 | 2.5 |
| Polyamine | 15.0 | 15.0 | 17.5 | 17.5 | 20.0 | 20.0 |
| Bayhibit AM | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 |
| C ₁₂ -C ₁₄ dimethyl hydroxyethyl quaternary ammonium chloride | - | 5.0 | 5.0 | - | - | - |
| Fabric softener active | - | - | 2.5 | 2.5 | - | - |
| Genamin C100 | 0.33 | - | 0.33 | 0.33 | 0.33 | - |
| Genapol V4463 | 0.2 | - | 0.2 | 0.2 | 0.2 | - |
| PARP3 | 2.0 | 1.0 | 0.2 | 0.5 | 0.1 | 0.16 |
| Water & minors | balance | balance | balance | balance | balance | balance |

Example 29

The following are non-limiting examples of odor-absorbing compositions suitable for spray-on applications:

10

| Examples | A | B | C | D | E |
|--------------------|-----------|-----------|-----------|-----------|-----------|
| Ingredients | Wt. % | Wt. % | Wt. % | Wt. % | Wt. % |
| HPBCD | 1.0 | - | 1.0 | - | 1.2 |
| RAMEB | - | 1.0 | - | 0.8 | - |
| Tetronic 901 | - | - | 0.1 | - | - |
| Silwet L-7604 | - | - | - | 0.1 | - |
| Silwet L-7600 | 0.1 | - | - | - | 0.1 |
| Bardac 2050 | - | - | - | 0.03 | - |
| Bardac 2250 | - | 0.2 | - | - | 0.1 |
| Diethylene glycol | - | 1.0 | - | - | 0.2 |
| Triethylene glycol | - | - | 0.1 | - | - |
| Ethanol | - | - | - | - | 2.5 |
| Perfume 1 | 0.1 | - | - | - | - |
| Perfume 2 | - | 0.05 | - | 0.1 | - |
| Perfume 3 | - | - | 0.1 | - | 0.1 |
| Kathon | 3 ppm | 3 ppm | 3 ppm | 3 ppm | - |
| HCl | to pH 4.5 | to pH 4.5 | to pH 3.5 | to pH 3.5 | to pH 3.5 |
| PARP2 | 5.0 | 2.0 | 1.0 | 0.2 | 0.16 |
| Distilled water | Bal. | Bal. | Bal. | Bal. | Bal. |

The perfume 1, 2, and 3 have the following compositions:

| Perfume | 1 | 2 | 3 |
|---------------------|-------|-------|-------|
| Perfume Ingredients | Wt. % | Wt. % | Wt. % |
| Anisic aldehyde | - | - | 2 |
| Benzophenone | 3 | 5 | - |
| Benzyl acetate | 10 | 15 | 5 |
| Benzyl salicylate | 5 | 20 | 5 |
| Cedrol | 2 | - | - |
| Citronelloi | 10 | - | 5 |
| Coumarin | - | - | 5 |
| Cymal | - | - | 3 |
| Dihydromyrcenol | 10 | - | 5 |

| | | | |
|--------------------------|-----|-----|-----|
| Flor acetate | 5 | - | 5 |
| Galaxolide | 10 | - | - |
| Lilial | 10 | 15 | 20 |
| Linalyl acetate | 4 | - | 5 |
| Linalool | 6 | 15 | 5 |
| Methyl dihydro jasmonate | 3 | 10 | 5 |
| Phenyl ethyl acetate | 2 | 5 | 1 |
| Phenyl ethyl alcohol | 15 | 15 | 20 |
| alpha-Terpineol | 5 | - | 8 |
| Vanillin | - | - | 1 |
| Total | 100 | 100 | 100 |

TOTAL 826.001

Claims

1-A process for making particles of amine reaction product of a compound containing a primary and/or secondary amine functional group and an active ketone or aldehyde containing component, and which comprises the steps of :

- a)-providing an amine reaction product, and
- b)-mixing therewith a carrier having a melting point of less than 30°C.

2-A process according to Claim 1, wherein the amine reaction product has a viscosity of higher than 1000cps.

3-A process according to either one of Claim 1 or 2, wherein the primary and/or secondary amine is selected from aminoaryl derivatives, polyamines, amino acids and derivatives thereof, substituted amines and amides, glucamines, dendrimers, polyvinylamines and derivatives thereof, and/or copolymer thereof, alkylene polyamine, polyaminoacid and copolymer thereof, cross-linked polyaminoacids, amino substituted polyvinylalcohol, polyoxyethylene bis amine or bis aminoalkyl, aminoalkyl piperazine and derivatives thereof, bis (amino alkyl) alkyl diamine linear or branched, and mixtures thereof; preferably selected from ethyl-4-amino benzoate, polyethyleneimine polymers; polylysine, cross-linked polylysine, N,N'-bis-(3-aminopropyl)-1,3-propanediamine linear or branched, 1,4-bis-(3-aminopropyl) piperazine, and mixtures thereof.

4-A process according to any one of Claims 1-3, wherein the active is selected from a flavour ketone or aldehyde ingredient, a pharmaceutical ketone or aldehyde active, a biocontrol ketone or aldehyde agent, a perfume ketone or aldehyde component, a refreshing cooling ketone or aldehyde agent and/or mixtures thereof.

5-A process according to any one of Claims 1-4, wherein said active component is an insect and/or moth repellent, preferably selected from citronellal, citral, N, N diethyl meta toluamide, Rotundial, 8-acetoxycarvotanacenone, and mixtures thereof.

6-A process according to any one of Claims 1-4, wherein said active component is an antimicrobial, preferably selected from Glutaraldehyde, Cinnamaldehyde, and mixtures thereof.

7-A process according to any one of Claims 1-4, wherein the active is a perfume, preferably selected from alpha-damascone, delta damascone, Carvone, Gamma-Methyl-Ionone; Damascenone, hedione, 2,4-dimethyl-3-cyclohexen-1-carboxaldehyde, Florhydral, Lillial, heliotropine, trans-2-nonenal, citral, and mixtures thereof.

8-A process according to any one of Claims 1-7, wherein the amount of amine reaction product ranges from 1 to 75%, preferably 5 to 30%, by weight of the processed amine reaction product.

9-A process according to any one of Claims 1-8, wherein the carrier has a melting point between minus 150°C and less than 30°C.

10-A process according to any one of Claims 1-9, wherein said carrier is a liquid carrier, preferably selected from hydrophobic and hydrophilic carrier, more preferably are selected from benzyl salicylate, diethyl phthalate, dipropylene glycol, methanol, ethanol, hydrophobic perfume composition, and mixtures thereof.

11-A process according to any one of Claims 1-10, wherein the amount of carrier material ranges from 3 to 95%, preferably from 15 to 80% and most preferably from 25 to 75%, by weight of the produced particles of the processed amine reaction product.

12-A process according to Claim 11 wherein the processed product is subsequently treated to form a coated particle.

13-A process according to any one of Claims 1-12, wherein the coating is made of water-soluble agglomerating agent.

14-A process according to Claim 13, wherein the water-soluble agglomerating agent is selected from water soluble organic polymeric compounds, water soluble

monomeric polycarboxylates, or their acid forms, homo or copolymeric polycarboxylic acids or their salts in which the polycarboxylic acid comprises at least two carboxylic radicals separated from each other by not more than two carbon atoms, carbonates, bicarbonates, borates, phosphates, sulfate salts, inorganic perhydrate salts, silicates, starch, cyclodextrin, and mixtures thereof, preferably selected from starch, carbonate, cyclodextrin, and mixtures thereof.

15-A process according to any one of Claims 1-14, wherein the amount of amine reaction product ranges from 1 to 75%, preferably 5 to 30%, by weight of the coated particle.

16-A processed amine reaction product as obtainable by the process of Claims 1-15.

17-A method of incorporating an amine reaction product according to Claim 16 into finished product, preferably by means selected from spraying, dry-addition, and mixtures thereof.

18-A finished composition comprising one or more laundry or cleaning ingredient and a processed amine reaction product according to Claim 16.

19-A composition according to Claim 18, wherein said composition is selected from a laundry composition, hard surface cleaning composition, and personal cleaning composition.

20-A method for delivering residual active to a surface which comprises the steps of contacting said surface with a processed product according to Claim 16, or composition according to either one of Claim 18 or 19, and thereafter contacting the treated surface with a material so that the active is released.

21- Use of a product as defined in Claim 16 for the manufacture of a laundry and/or cleaning composition for delivering residual fragrance on a surface on which it is applied.

22-Use of a product as defined in Claim 16 for the manufacture of a laundry and/or cleaning composition for delivering residual fragrance and fabric care onto the fabrics on which it is applied.

23-A packaged composition comprising the processed product of Claim 16 or composition of either one of Claim 18 or 19.

1004943 123401
TOTAL 826007

DECLARATION COMBINED WITH POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

"Process for Producing Particles of Amine Reaction Product"

bearing the above listed Procter & Gamble Company Case number, the specification of which was filed as PCT/US00/18468, designating at least the United States of America, with the United States Receiving Office on 06 July 2000.

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37 Code of Federal Regulations §1.56.

I hereby claim foreign priority benefits under Title 35 United States Code §119 of any foreign application(s) for patent or inventor's certificate or of any PCT international application(s) designating at least one country other than the United States of America listed below and have also identified below any foreign application for patent or Inventor's certificate having a filing date before that of the application on which priority is claimed:

PRIOR FOREIGN APPLICATION(S) TO WHICH WE CLAIM PRIORITY:

99870146.0 EPO 08 July 1999

I hereby claim the benefit under Title 35 United States Code §120 of any United States application(s) or PCT international application(s) designating the United States of America that is/are listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35 United States Code §112, I acknowledge the duty to disclose material information as defined in Title 37 Code of Federal Regulations §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

| | | |
|---------------------|---------------|--|
| (Appln. Serial No.) | (Filing Date) | (Status)(patented, pending, abandoned) |
| (Appln. Serial No.) | (Filing Date) | (Status)(patented, pending, abandoned) |

1001928.12001

I hereby appoint the following as my attorney(s) or agent(s) with full power of substitution to prosecute this application and transact all business in the Patent and Trademark office connected therewith:

| Name | Registration No. | Associate Power of Attorney Attached [] Yes [] No |
|------|------------------|---|
|------|------------------|---|

| | | |
|-------------------|--------|--|
| Jacobus C. Rasser | 37,043 | |
| Donald E. Hasse | 29,387 | |
| T. David Reed | 32,931 | |
| Eileen L. Hughett | 34,352 | |
| Timothy B. Guffey | 41,048 | |
| Emelyn L. Hiland | 41,501 | |

SEND CORRESPONDENCE TO:

| | | | |
|---|------------|-------|----------------|
| T. David Reed, c/o The Procter & Gamble Company | | | (513) 627-7025 |
| Name | | | Phone No. |
| 5299 Spring Grove Avenue | Cincinnati | Ohio | 45217-1087 |
| Street | City | State | Zip Code |

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willfull false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such will ful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00
Full Name of sole or first inventor: BIJSCH, Alfred (NMN)
Inventor's signature [Signature] Date: July 17, 2000
Residence: Handelsstraat 210, B-1840 Londerzeel, BEX BE
Citizenship: German
Post Office Address: Procter & Gamble Eurocor N.V.
100 Temselaan, B-1853 Strombeek-Bever, Belgium

2-00
Full Name of second joint inventor, if any: HOMBLÉ, Marcel (NMN)
Inventor's signature [Signature] Date: July 17, 2000
Residence: Kouter 5, B-3060 Bertem, BEX BE
Citizenship: Belgian
Post Office Address: Procter & Gamble Eurocor N.V.
100 Temselaan, B-1853 Strombeek-Bever, Belgium

3-00
Full name of third joint inventor, if any: LAUDAMIEL, Christophe (NMN)
Inventor's signature [Signature] Date: July 17, 2000
Residence: Rue de la Violette 20A, B-1000 Brussels, BEX BE
Citizenship: French
Post Office Address: Procter & Gamble Eurocor N.V.
100 Temselaan, B-1853 Strombeek-Bever, Belgium

4-00
Full Name of fourth joint inventor, if any: SMEETS, Johan (NMN)
Inventor's signature [Signature] Date: July 25, 2000
Residence: Bollenberg 79, B-3210 Lubbeek, BEX BE
Citizenship: Belgian
Post Office Address: Procter & Gamble Eurocor N.V.
100 Temselaan, B-1853 Strombeek-Bever, Belgium

5-00
Full name of fifth joint inventor, if any:

TRUJILLO, Rafael (NMN)

Inventor's signature

Date: July 17, 2000

Residence:

Avenue Princesse Paolo 39, B-1410 Waterloo, BEX

BE

Citizenship:

Mexican

Post Office Address:

Procter & Gamble Eurocor N.V.

100 Temselaan, B-1853 Strombeek-Bever, Belgium

6-00
Full Name of sixth joint inventor, if any:

WEVERS, Jean (NMN)

Inventor's signature

Date: July 17, 2000

Residence:

Heide 17, B-1840 Steenhuffel, BEX

BE

Citizenship:

Belgian

Post Office Address:

Procter & Gamble Eurocor N.V.

100 Temselaan, B-1853 Strombeek-Bever, Belgium

10019178 199104